

Product datasheet for RC200455L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

LOXL2 (NM_002318) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: LOXL2 (NM_002318) Human Tagged ORF Clone Lentiviral Particle

Symbol: LOXL2

Synonyms: LOR; LOR2; WS9-14

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM_002318

 ORF Size:
 2322 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC200455).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 002318.2</u>

 RefSeq Size:
 3810 bp

 RefSeq ORF:
 2325 bp

 Locus ID:
 4017

 UniProt ID:
 Q9Y4K0

 Cytogenetics:
 8p21.3

Domains: SR, Lysyl_oxidase

Protein Families: Druggable Genome, Secreted Protein





ORIGENE

MW: 86.7 kDa

Gene Summary:

This gene encodes a member of the lysyl oxidase gene family. The prototypic member of the family is essential to the biogenesis of connective tissue, encoding an extracellular copperdependent amine oxidase that catalyses the first step in the formation of crosslinks in collagens and elastin. A highly conserved amino acid sequence at the C-terminus end appears to be sufficient for amine oxidase activity, suggesting that each family member may retain this function. The N-terminus is poorly conserved and may impart additional roles in developmental regulation, senescence, tumor suppression, cell growth control, and chemotaxis to each member of the family. [provided by RefSeq, Jul 2008]