

Product datasheet for RC200464L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Hexokinase 1 (HK1) (NM 000188) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Hexokinase 1 (HK1) (NM_000188) Human Tagged ORF Clone Lentiviral Particle

Symbol:

hexokinase; HK; HK1-ta; HK1-tb; HK1-tc; HKD; HKI; HMSNR; HXK1; NEDVIBA; RP79 Synonyms:

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK NM 000188 ACCN:

ORF Size: 2751 bp

ORF Nucleotide

OTI Disclaimer:

Sequence:

The ORF insert of this clone is exactly the same as(RC200464).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 000188.1

RefSeq Size: 3580 bp RefSeq ORF: 2754 bp Locus ID: 3098

UniProt ID: P19367

Cytogenetics: 10q22.1

Domains: hexokinase

Protein Families: Druggable Genome





Hexokinase 1 (HK1) (NM_000188) Human Tagged ORF Clone Lentiviral Particle - RC200464L3V

Protein Pathways: Amino sugar and nucleotide sugar metabolism, Fructose and mannose metabolism,

Galactose metabolism, Glycolysis / Gluconeogenesis, Insulin signaling pathway, Metabolic

pathways, Starch and sucrose metabolism, Type II diabetes mellitus

MW: 102.3 kDa

Gene Summary: Hexokinases phosphorylate glucose to produce glucose-6-phosphate, the first step in most

glucose metabolism pathways. This gene encodes a ubiquitous form of hexokinase which localizes to the outer membrane of mitochondria. Mutations in this gene have been

associated with hemolytic anemia due to hexokinase deficiency. Alternative splicing of this gene results in several transcript variants which encode different isoforms, some of which

are tissue-specific. [provided by RefSeq, Apr 2016]