

## Product datasheet for RC200472L3

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Glucokinase (GCK) (NM\_000162) Human Tagged Lenti ORF Clone

#### **Product data:**

**Product Type:** Expression Plasmids

Product Name: Glucokinase (GCK) (NM\_000162) Human Tagged Lenti ORF Clone

Tag: Myc-DDK

Symbol: Glucokinase

Synonyms: FGQTL3; GK; GLK; HHF3; HK4; HKIV; HXKP; LGLK; MODY2; PNDM1

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

E. coli Selection: Chloramphenicol (34 ug/mL)


ORF Nucleotide The ORF insert of this cl


Sequence:

The ORF insert of this clone is exactly the same as(RC200472).

**Restriction Sites:** Sgfl-Mlul

**Cloning Scheme:** 





<sup>\*</sup> The last codon before the Stop codon of the ORF.

**ACCN:** NM\_000162

ORF Size: 1395 bp



### Glucokinase (GCK) (NM\_000162) Human Tagged Lenti ORF Clone - RC200472L3

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**RefSeq:** <u>NM 000162.2</u>

 RefSeq Size:
 2741 bp

 RefSeq ORF:
 1398 bp

 Locus ID:
 2645

 UniProt ID:
 P35557

**Cytogenetics:** 7p13

**Protein Families:** Druggable Genome

**Protein Pathways:** Amino sugar and nucleotide sugar metabolism, Galactose metabolism, Glycolysis /

Gluconeogenesis, Insulin signaling pathway, Maturity onset diabetes of the young, Metabolic

pathways, Starch and sucrose metabolism, Type II diabetes mellitus

**MW:** 52.2 kDa


**Gene Summary:** This gene encodes a member of the hexokinase family of proteins. Hexokinases

phosphorylate glucose to produce glucose-6-phosphate, the first step in most glucose metabolism pathways. In contrast to other forms of hexokinase, this enzyme is not inhibited by its product glucose-6-phosphate but remains active while glucose is abundant. The use of multiple promoters and alternative splicing of this gene result in distinct protein isoforms that exhibit tissue-specific expression in the pancreas and liver. In the pancreas, this enzyme plays a role in glucose-stimulated insulin secretion, while in the liver, this enzyme is important in glucose uptake and conversion to glycogen. Mutations in this gene that alter enzyme

activity have been associated with multiple types of diabetes and hyperinsulinemic hypoglycemia. [provided by RefSeq, Aug 2017]



# **Product images:**



Double digestion of RC200472L3 using Sgfl and Mlul  $\,$