

Product datasheet for RC200485L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CPT1A (NM_001031847) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: CPT1A (NM_001031847) Human Tagged ORF Clone Lentiviral Particle

Symbol: CPT1A

Synonyms: CPT1; CPT1-L; L-CPT1

Mammalian Cell

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK

ACCN: NM_001031847

ORF Size: 2268 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC200485).

Sequence:

Cytogenetics:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 001031847.1

 RefSeq Size:
 2671 bp

 RefSeq ORF:
 2271 bp

 Locus ID:
 1374

 UniProt ID:
 P50416

Protein Families: Druggable Genome, Transmembrane

11q13.3

Protein Pathways: Adipocytokine signaling pathway, Fatty acid metabolism, PPAR signaling pathway





ORIGENE

MW: 86.2 kDa

Gene Summary:

The mitochondrial oxidation of long-chain fatty acids is initiated by the sequential action of carnitine palmitoyltransferase I (which is located in the outer membrane and is detergent-labile) and carnitine palmitoyltransferase II (which is located in the inner membrane and is detergent-stable), together with a carnitine-acylcarnitine translocase. CPT I is the key enzyme in the carnitine-dependent transport across the mitochondrial inner membrane and its deficiency results in a decreased rate of fatty acid beta-oxidation. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]