

Product datasheet for RC200494L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CDK2 (NM 001798) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: CDK2 (NM_001798) Human Tagged ORF Clone Lentiviral Particle

Symbol:

CDKN2; p33(CDK2) Synonyms:

Mammalian Cell

Selection:

ACCN:

None

Vector: pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK NM 001798

ORF Size: 894 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC200494).

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of OTI Disclaimer:

> reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 001798.2

RefSeq Size: 2301 bp RefSeq ORF: 897 bp Locus ID: 1017 **UniProt ID:** P24941 Cytogenetics: 12q13.2

Domains: pkinase, TyrKc, S_TKc

Protein Families: Druggable Genome, Protein Kinase



CDK2 (NM_001798) Human Tagged ORF Clone Lentiviral Particle - RC200494L1V

Protein Pathways: Cell cycle, Oocyte meiosis, p53 signaling pathway, Pathways in cancer, Progesterone-

mediated oocyte maturation, Prostate cancer, Small cell lung cancer

MW: 33.9 kDa

Gene Summary: This gene encodes a member of a family of serine/threonine protein kinases that participate

in cell cycle regulation. The encoded protein is the catalytic subunit of the cyclin-dependent protein kinase complex, which regulates progression through the cell cycle. Activity of this protein is especially critical during the G1 to S phase transition. This protein associates with and regulated by other subunits of the complex including cyclin A or E, CDK inhibitor p21Cip1 (CDKN1A), and p27Kip1 (CDKN1B). Alternative splicing results in multiple transcript variants.

[provided by RefSeq, Mar 2014]