

Product datasheet for RC200513L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

HAND1 (NM_004821) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: HAND1 (NM 004821) Human Tagged ORF Clone Lentiviral Particle

Symbol: HAND1

Synonyms: bHLHa27; eHand; Hxt; Thing1

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_004821

ORF Size: 645 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC200513).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 004821.1

 RefSeq Size:
 1750 bp

 RefSeq ORF:
 648 bp

 Locus ID:
 9421

 UniProt ID:
 096004

Cytogenetics: 5q33.2

Protein Families: Druggable Genome, ES

Protein Families: Druggable Genome, ES Cell Differentiation/IPS, Transcription Factors

MW: 23.4 kDa







Gene Summary:

The protein encoded by this gene belongs to the basic helix-loop-helix family of transcription factors. This gene product is one of two closely related family members, the HAND proteins, which are asymmetrically expressed in the developing ventricular chambers and play an essential role in cardiac morphogenesis. Working in a complementary fashion, they function in the formation of the right ventricle and aortic arch arteries, implicating them as mediators of congenital heart disease. In addition, it has been suggested that this transcription factor may be required for early trophoblast differentiation. [provided by RefSeq, Jul 2008]