

Product datasheet for **RC200523**

HSPA2 (NM_021979) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	HSPA2 (NM_021979) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	HSPA2
Synonyms:	HSP70-2; HSP70-3
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

**ORF Nucleotide
Sequence:**

>RC200523 ORF sequence
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGGATCGCC**

ATGCTGCCCGTGGCCCGGCTATCGGCATCGACCTGGGCACCACCTATTCTGTCGTCGGGGTCTTCCAAC
 ATGGCAAGGTGGAGATCATCGCCAACGACCAGGGCAATCGCACCACCCAGCTACGTGGCCTTCACGGA
 CACCGAGCGCTCATCGGCGACGCCAAGAACCAGGTGGCCATGAACCCACCAACACCATCTTCGAC
 GCCAAGAGGCTGATTGGACGAAATTCGAGGATGCCACAGTGCAGTCGGATATGAAACACTGGCCGTTC
 GGGTGGTGGAGGAGGCAAGCCAAAGTGAAGTAGAGTACAAGGGGAGACCAAGACCTTCTTCCC
 AGAGGAGATATCCTCCATGGTCTCACGAAGATGAAGGAGATCGCGGAAGCCTACCTGGGGGCAAGGTG
 CACAGCGCGTACATAACGGTCCCGCCTATTTCAACGACTCGCAGCGCCAGGCCACCAAGGACGCAGGCA
 CCATCACGGGGCTCAATGTGCTGCGCATCATCAACGAGCCACGGCGGCCCATCGCCTACGGCCTGGA
 CAAGAAGGGCTGCGCGGGCGGCGAGAAGAAGTGTCTCATCTTTGACCTGGGCGGTGGCACTTTCGACGTG
 TCCATCTGACCATCGAGGATGGCATCTTCGAGGTGAAGTCCACGGCCGGCGACACCCACCTGGGCGGTG
 AGGACTTCGACAACCGCATGGTGAAGCCACTGGCGGAGGAGTTCAAGCGCAAGCACAAGAAGGACATTGG
 GCCCAACAAGCGCGCCGTGAGGCGGCTGCGCACCGCTTGCAGCGCGCCAAGCGCACCTGAGCTCGTCC
 ACGCAGGCGAGCATCGAGATCGACTCGCTTACGAGGGCGTGGACTTCTATACGTCCATCACGCGCGCC
 GCTTCGAGGAGCTCAATGCCGACCTCTTTCGCGGGACCTGGAGCCGGTGGAGAAGGCGCTGCGCGACGC
 CAAGCTGGACAAGGGCCAGATCCAGGAGATCGTGTGGTGGGCGGCTCCACTCGTATCCCAAGATCCAG
 AAGCTGCTGCAGGATTTCTTCAACGGCAAGGAGCTGAACAAGAGCATCAACCCGACGAGGCGGTGGCCT
 ATGGCGCCGCGGTGCAGGCGGCCATCCTCATCGGCGACAATCAGAGAATGTGCAGGACCTGCTGCTACT
 CGACGTGACCCCGTTGTCGCTGGGCATCGAGACAGCTGGCGGTGTCATGACCCCACTCATCAAGAGGAAC
 ACCACGATCCCCACCAAGCAGACGACGACCTTACCACCTACTCGGACAACCAGAGCAGCGTACTGGTGC
 AGGTATACGAGGGCGAACGGGCCATGACCAAGGACAATAACCTGCTGGGCAAGTTCGACCTGACCGGGAT
 TCCCCCTGCGCCTCGCGGGTCCCCAAATCGAGGTTACCTTCGACATTGACGCAATGGCATCCTTAAC
 GTTACCGCCGCGACAAGAGCACCGGTAAGGAAAAAAAATCACCATACCAATGACAAAGGTGCTCTGA
 GCAAGGACGACATTGACCGGATGGTGCAGGAGGCGGAGCGGTACAATCGGAAGATGAGGCGAATCGCGA
 CCGAGTCGCGGCAAAAACGCCCTGGAGTCTATACTACAACATCAAGCAGACGGTGAAGACGAGAAA
 CTGAGGGGCAAGATTAGCGAGCAGGACAAAAACAAGATCCTCGACAAGTGTGAGGAGGTGATCAACTGGC
 TCGACCGAAACCAGATGGCAGAGAAAGATGAGTATGAACACAAGCAGAAAGAGCTCGAAAGAGTTTGCAA
 CCCCATCATCAGCAAACCTTTACCAAGGTGGTCTGGCGGCGGAGCGGCGGCGGCGGTTACGAGACCTCC
 GGGGACCCACCATCGAAGAAGTGGAC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC200523 protein sequence
 Red=Cloning site Green=Tags(s)


MSARGPAIGIDLGTTYSCVGVFQHGKVEIIANDQGNRTTPSYVAFDTERLIGDAAKNQVAMNPTNTIFD
 AKRLIGRKFEDATVQSDMKHWPFRVVSSEGGKPKVQVEYKGETKTFPPEEISSMVLTKMKEIAEAYLGGKV
 HSAVITVPAYFNDSQRQATKDAGTITGLNVLRIINEPTAAAIAYGLDKKGCAGGEKNVIFDLGGGTFDV
 SILTIEDGIFEVKSTAGDTHLGGEDFDNRMVSHLAEFEFKRKHKKDIGPNKRAVRRLRTACERAKRTLSSS
 TQASIEIDSLYEGVDFYTSITRARFEELNADLFRGTLEPVEKALRDAKLDKGQIQEIVLVGGSTRIPKIQ
 KLLQDFFNKELNKSINPDEAVAYGAAVQAAAILIGDKSENVQDLLLDDVTPLSLGIETAGGYMTPLIKRN
 TTIPTKQTQTFTTYSNQS SVLVQVYEGERAMTKDNNLLGKFDLTGIPPAPRGVPIEVTFDIDANGILN
 VTAADKSTGKENKIIITNDKGRLSKDDIDRMVQEAERYKSEDEANRDRVAAKNALESYTYNIKQTVDEK
 LRGKISEQDNKILDKCQEVINWLDNRNQMAEKDEYEHKQKELERVCPNPIISKLYQGGPGGGGGGGSGAS
 GGPTIEEVD

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mk6149_h04.zip

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_021979

ORF Size: 1917 bp

OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_021979.3](#)

RefSeq Size: 2802 bp

RefSeq ORF: 1920 bp

Locus ID: 3306

UniProt ID: [P54652](#)

Cytogenetics: 14q23.3

Domains: HSP70


Protein Families: Stem cell - Pluripotency

Protein Pathways: Antigen processing and presentation, Endocytosis, MAPK signaling pathway, Spliceosome


MW: 70 kDa

Gene Summary:


Molecular chaperone implicated in a wide variety of cellular processes, including protection of the proteome from stress, folding and transport of newly synthesized polypeptides, activation of proteolysis of misfolded proteins and the formation and dissociation of protein complexes. Plays a pivotal role in the protein quality control system, ensuring the correct folding of proteins, the re-folding of misfolded proteins and controlling the targeting of proteins for subsequent degradation. This is achieved through cycles of ATP binding, ATP hydrolysis and ADP release, mediated by co-chaperones. The affinity for polypeptides is regulated by its nucleotide bound state. In the ATP-bound form, it has a low affinity for substrate proteins. However, upon hydrolysis of the ATP to ADP, it undergoes a conformational change that increases its affinity for substrate proteins. It goes through repeated cycles of ATP hydrolysis and nucleotide exchange, which permits cycles of substrate binding and release (PubMed:26865365). Plays a role in spermatogenesis. In association with SHCBP1L may participate in the maintenance of spindle integrity during meiosis in male germ cells (By similarity).[UniProtKB/Swiss-Prot Function]

Product images:


Circular map for RC200523



Western blot validation of overexpression lysate (Cat# [LY402898]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC200523 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).



Coomassie blue staining of purified HSPA2 protein (Cat# [TP300523]). The protein was produced from HEK293T cells transfected with HSPA2 cDNA clone (Cat# RC200523) using MegaTran 2.0 (Cat# [TT210002]).