

Product datasheet for RC200539L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Calpain 6 (CAPN6) (NM 014289) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Calpain 6 (CAPN6) (NM_014289) Human Tagged ORF Clone Lentiviral Particle

Symbol: Calpain 6

Synonyms: CalpM; CANPX; CAPNX; DJ914P14.1

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM_014289

ORF Size: 1923 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC200539).

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 014289.2

RefSeq Size:3604 bpRefSeq ORF:1926 bp

Locus ID: 827

UniProt ID: Q9Y6Q1

Cytogenetics: Xq23

Domains: C2, Calpain_III

Protein Families: Druggable Genome, Protease





ORIGENE

MW: 74.6 kDa

Gene Summary:

Calpains are ubiquitous, well-conserved family of calcium-dependent, cysteine proteases. The calpain proteins are heterodimers consisting of an invariant small subunit and variable large subunits. The large subunit possesses a cysteine protease domain, and both subunits possess calcium-binding domains. Calpains have been implicated in neurodegenerative processes, as their activation can be triggered by calcium influx and oxidative stress. The protein encoded by this gene is highly expressed in the placenta. Its C-terminal region lacks any homology to the calmodulin-like domain of other calpains. The protein lacks critical active site residues and thus is suggested to be proteolytically inactive. The protein may play a role in tumor formation by inhibiting apoptosis and promoting angiogenesis. [provided by RefSeq, Nov 2009]