

Product datasheet for RC200568L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Integrin beta 4 binding protein (EIF6) (NM_002212) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Integrin beta 4 binding protein (EIF6) (NM_002212) Human Tagged ORF Clone Lentiviral

Particle

Symbol: Integrin beta 4 binding protein

Synonyms: b(2)gcn; CAB; eIF-6; EIF3A; ITGB4BP; p27(BBP); p27BBP

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_002212

ORF Size: 735 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC200568).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 002212.2</u>

 RefSeq Size:
 1135 bp

 RefSeq ORF:
 738 bp

 Locus ID:
 3692

 UniProt ID:
 P56537

Cytogenetics: 20q11.22

Domains: eIF6





Integrin beta 4 binding protein (EIF6) (NM_002212) Human Tagged ORF Clone Lentiviral Particle – RC200568L4V

Protein Families: Druggable Genome

MW: 26.6 kDa

Gene Summary: Hemidesmosomes are structures which link the basal lamina to the intermediate filament

cytoskeleton. An important functional component of hemidesmosomes is the integrin beta-4 subunit (ITGB4), a protein containing two fibronectin type III domains. The protein encoded by this gene binds to the fibronectin type III domains of ITGB4 and may help link ITGB4 to the intermediate filament cytoskeleton. The encoded protein, which is insoluble and found both in the nucleus and in the cytoplasm, can function as a translation initiation factor and prevent the association of the 40S and 60S ribosomal subunits. Multiple non-protein coding transcript

variants and variants encoding two different isoforms have been found for this gene.

[provided by RefSeq, Jun 2012]