

## Product datasheet for RC200601L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## HIV TAT specific factor 1 (HTATSF1) (NM\_014500) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** HIV TAT specific factor 1 (HTATSF1) (NM\_014500) Human Tagged ORF Clone Lentiviral Particle

Symbol: HIV TAT specific factor 1

**Synonyms:** dJ196E23.2; TAT-SF1; TATSF1

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_014500

ORF Size: 2265 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC200601).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** <u>NM 014500.4</u>

 RefSeq Size:
 2788 bp

 RefSeq ORF:
 2268 bp

 Locus ID:
 27336

 UniProt ID:
 043719

 Cytogenetics:
 Xq26.3

 Domains:
 RRM





## HIV TAT specific factor 1 (HTATSF1) (NM\_014500) Human Tagged ORF Clone Lentiviral Particle – RC200601L4V

**Protein Families:** Transcription Factors

MW: 86.3 kDa

**Gene Summary:** The protein encoded by this gene functions as a cofactor for the stimulation of

transcriptional elongation by HIV-1 Tat, which binds to the HIV-1 promoter through Tat-TAR interaction. This protein may also serve as a dual-function factor to couple transcription and splicing and to facilitate their reciprocal activation. Alternatively spliced transcript variants

have been found for this gene.[provided by RefSeq, Sep 2009]