

# Product datasheet for RC200606L2

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## SAPK4 (MAPK13) (NM\_002754) Human Tagged Lenti ORF Clone

#### **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** SAPK4 (MAPK13) (NM\_002754) Human Tagged Lenti ORF Clone

Tag: mGFP Symbol: SAPK4

Synonyms: MAPK-13; MAPK 13; p38delta; PRKM13; SAPK4

Mammalian Cell None

Selection:

Vector:pLenti-C-mGFP (PS100071)E. coli Selection:Chloramphenicol (34 ug/ml

E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this clone

Sequence:

The ORF insert of this clone is exactly the same as(RC200606).

**Restriction Sites:** Sgfl-Mlul

**Cloning Scheme:** 





<sup>\*</sup> The last codon before the Stop codon of the ORF.

**ACCN:** NM\_002754

ORF Size: 1095 bp



## SAPK4 (MAPK13) (NM\_002754) Human Tagged Lenti ORF Clone - RC200606L2

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**RefSeq:** NM 002754.3, NP 002745.1

 RefSeq Size:
 6348 bp

 RefSeq ORF:
 1098 bp

 Locus ID:
 5603

 UniProt ID:
 015264

Cytogenetics: 6p21.31

**Domains:** pkinase, TyrKc, S\_TKc

**Protein Families:** Druggable Genome, Protein Kinase

**Protein Pathways:** Amyotrophic lateral sclerosis (ALS), Epithelial cell signaling in Helicobacter pylori infection, Fc

epsilon RI signaling pathway, GnRH signaling pathway, Leukocyte transendothelial migration, MAPK signaling pathway, Neurotrophin signaling pathway, NOD-like receptor signaling pathway, Progesterone-mediated oocyte maturation, RIG-I-like receptor signaling pathway, T cell receptor signaling pathway, Toll-like receptor signaling pathway, VEGF signaling pathway

**MW:** 42.1 kDa


**Gene Summary:** This gene encodes a member of the mitogen-activated protein (MAP) kinase family. MAP

kinases act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. The encoded protein is a p38 MAP kinase and is activated by proinflammatory cytokines and cellular stress. Substrates of the encoded protein include the transcription factor ATF2 and the microtubule dynamics regulator stathmin. Alternatively spliced transcript

variants have been observed for this gene. [provided by RefSeq, Jul 2012]



# **Product images:**



Circular map for RC200606L2