

Product datasheet for **RC200615**

MEST (NM_002402) Human Tagged ORF Clone

Product data:

| | |
|---------------------------|---|
| Product Type: | Expression Plasmids |
| Product Name: | MEST (NM_002402) Human Tagged ORF Clone |
| Tag: | Myc-DDK |
| Symbol: | MEST |
| Synonyms: | PEG1 |
| Mammalian Cell Selection: | Neomycin |
| Vector: | pCMV6-Entry (PS100001) |
| E. coli Selection: | Kanamycin (25 ug/mL) |
| ORF Nucleotide Sequence: | >RC200615 ORF sequence Red=Cloning site Blue=ORF Green=Tags(s) |

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGGTGC GCCGAGATCGCTCCGCAGGATGAGGGAGTGGTGGTCCAGGTGGGGCTGCTGGCCGTGCCCC
TGCTTGCTGCGTACCTGCACATCCACCCCTCAGCTCTCCCCTGCCCTCACTCATGGAAGTCTTCAGG
CAAGTTTTCACTTACAAGGGACTGCGTATCTTACCAAGACTCTGTGGGTGTGGTTGGAAGTCCAGAG
ATAGTTGTGCTTTACACGGTTTTCCAACATCCAGCTACGACTGGTACAAGATTTGGGAAGTCTGACCT
TGAGGTTTCATCGGGTATTGCCCTTGATTTCTTAGGCTTTGGCTTCAGTGACAAACCGAGACCACATCA
CTATTCATATTTGAGCAGGCCAGCATCGTGGAAGCGCTTTTGCGGCATCTGGGGCTCCAGAACCGCAGG
ATCAACCTTTCTCATGACTATGGAGATATTGTTGCTCAGGAGCTTCTCTACAGGTACAAGCAGAATC
GATCTGGTCCGGCTTACCATAAAGAGTCTCTGTCTGCAATGGAGGTATCTTTCTGAGACTCACCGTCC
ACTCCTTCTCCAAAAGCTACTCAAAGATGGAGGTGTGCTGTACCCATCCTCACACGACTGATGAACTTC
TTTGTATTCTCTCGAGGTCTCACCCAGTCTTTGGGCCGTATACTCGGCCCTCTGAGAGTGTGGG
ACATGTGGGCAGGGATCCGCAACAATGACGGGAAGTATGTCATTGACAGTCTCTTACAGTACATCAATCA
GAGGAAGAAGTTCAGAAGCGCTGGGTGGGAGCTCTTGCCTCTGTAAGTATCCCCATTATTTTATCTAT
GGCCATTGGATCCTGTAATCCCTATCCAGAGTTTTGGAGCTGTACAGGAAAACGCTGCCGCGGTCCA
CAGTGTGATTCTGGATGACCACATTAGCCACTATCCACAGCTAGAGGATCCCATGGGCTCTTTGAATGC
ATATATGGGCTTCATCAACTCTTC

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)

Protein Sequence: >RC200615 protein sequence
Red=Cloning site Green=Tags(s)

```
MVRRDRLRRMREWWVQVGLLAVPLLAAYLHIPPQLSPALHSWKSSGKFFTYKGLRIFYQDSVGVGSPE
IVVLLHGFPTSSYDWYKIWEGLTLRFHRVIALDFLGFGFSDKPRPHHYSIFEQASIVEALLRHLGLQNR
INLLSHDYGDIVAQELLRYKQNRSGRLTIKSLCLNNGGIFPETHRPLLKLLKDGVLSPILTRLMNF
FVFSRGLTPVFGPYTRPSESELWDMWAGIRNNDGNLVIDSLLQYINQRKFRRRVWVGALASVTIPIHFIY
GPLDPVNPYEFLELYRKTLPRSTVSIILDDHISHYPQLEDPMGFLNAYMGFINSF
```

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mk6271_f01.zip

Restriction Sites: SgfI-MluI

Cloning Scheme:

Cloning sites used for ORF Shutting:



* The last codon before the Stop codon of the ORF

ACCN: NM_002402

ORF Size: 1005 bp


OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.


Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

| | |
|-------------------------------|--|
| Reconstitution Method: | <ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C. |
| RefSeq: | NM_002402.4 |
| RefSeq Size: | 2513 bp |
| RefSeq ORF: | 1008 bp |
| Locus ID: | 4232 |
| UniProt ID: | Q5EB52 |
| Cytogenetics: | 7q32.2 |
| Domains: | abhydrolase |
| Protein Families: | Protease, Transmembrane |
| MW: | 38.8 kDa |
| Gene Summary: | <p>This gene encodes a member of the alpha/beta hydrolase superfamily. It is imprinted, exhibiting preferential expression from the paternal allele in fetal tissues, and isoform-specific imprinting in lymphocytes. The loss of imprinting of this gene has been linked to certain types of cancer and may be due to promotor switching. The encoded protein may play a role in development. Alternatively spliced transcript variants encoding multiple isoforms have been identified for this gene. Pseudogenes of this gene are located on the short arm of chromosomes 3 and 4, and the long arm of chromosomes 6 and 15. [provided by RefSeq, Dec 2011]</p> |

Product images:



Circular map for RC200615



Western blot validation of overexpression lysate (Cat# [LY419353]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC200615 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).