

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC200621L1V

Epoxide hydrolase (EPHX1) (NM_000120) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Epoxide hydrolase (EPHX1) (NM_000120) Human Tagged ORF Clone Lentiviral Particle
Symbol:	EPHX1
Synonyms:	EPHX; EPOX; HYL1; MEH
Mammalian Cell Selection:	None
Vector:	pLenti-C-Myc-DDK (PS100064)
Tag:	Myc-DDK
ACCN:	NM_000120
ORF Size:	1365 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC200621).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 000120.2</u>
RefSeq Size:	1847 bp
RefSeq ORF:	1368 bp
Locus ID:	2052
UniProt ID:	<u>P07099</u>
Cytogenetics:	1q42.12
Domains:	abhydrolase
Protein Families:	Druggable Genome, Protease



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Epoxide hydrolase (EPHX1) (NM_000120) Human Tagged ORF Clone Lentiviral Particle – RC200621L1V	
Protein Pathwa	: Metabolism of xenobiotics by cytochrome P450
MW:	52.9 kDa
Gene Summary	Epoxide hydrolase is a critical biotransformation enzyme that converts epoxides from the degradation of aromatic compounds to trans-dihydrodiols which can be conjugated and excreted from the body. Epoxide hydrolase functions in both the activation and detoxification of epoxides. Mutations in this gene cause preeclampsia, epoxide hydrolase deficiency or increased epoxide hydrolase activity. Alternatively spliced transcript variants encoding the same protein have been found for this gene.[provided by RefSeq, Dec 2008]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US