

## Product datasheet for RC200629L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **UBE3A (NM\_130838) Human Tagged ORF Clone Lentiviral Particle**

**Product data:** 

**Product Type:** Lentiviral Particles

Product Name: UBE3A (NM\_130838) Human Tagged ORF Clone Lentiviral Particle

Symbol: UBE3A

Synonyms: ANCR; AS; E6-AP; EPVE6AP; HPVE6A

**Mammalian Cell** 

Selection:

ACCN:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

NM 130838

Tag: Myc-DDK

ORF Size: 2556 bp

**ORF Nucleotide** 

Sequence:

**UniProt ID:** 

The ORF insert of this clone is exactly the same as(RC200629).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** <u>NM 130838.1</u>

RefSeq Size:4491 bpRefSeq ORF:2559 bpLocus ID:7337

Cytogenetics: 15q11.2

Domains: HECT

**Protein Families:** Druggable Genome

Q05086





## UBE3A (NM\_130838) Human Tagged ORF Clone Lentiviral Particle - RC200629L3V

**Protein Pathways:** Ubiquitin mediated proteolysis

**MW:** 97.8 kDa

**Gene Summary:** This gene encodes an E3 ubiquitin-protein ligase, part of the ubiquitin protein degradation

system. This imprinted gene is maternally expressed in brain and biallelically expressed in other tissues. Maternally inherited deletion of this gene causes Angelman Syndrome, characterized by severe motor and intellectual retardation, ataxia, hypotonia, epilepsy, absence of speech, and characteristic facies. The protein also interacts with the E6 protein of human papillomavirus types 16 and 18, resulting in ubiquitination and proteolysis of tumor protein p53. Alternative splicing of this gene results in three transcript variants encoding three isoforms with different N-termini. Additional transcript variants have been described, but their full length nature has not been determined. [provided by RefSeq, Jul 2008]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US