

## **Product datasheet for RC200633**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Sumo 1 (SUMO1) (NM\_003352) Human Tagged ORF Clone

#### **Product data:**

**Product Type:** Expression Plasmids

Tag: Myc-DDK
Symbol: Sumo 1

Synonyms: DAP1; GMP1; OFC10; PIC1; SENP2; SMT3; SMT3C; SMT3H3; UBL1

Mammalian Cell Neomycin

Selection:

Vector:

pCMV6-Entry (PS100001)


E. coli Selection: Kanamycin (25 ug/mL)

Chromatograms: <a href="https://cdn.origene.com/chromatograms/mk6408">https://cdn.origene.com/chromatograms/mk6408</a> b07.zip

**Restriction Sites:** Sgfl-Mlul

Cloning Scheme:





<sup>\*</sup> The last codon before the Stop codon of the ORF

**ACCN:** NM\_003352

ORF Size: 303 bp

### Sumo 1 (SUMO1) (NM\_003352) Human Tagged ORF Clone | RC200633

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**Note:** Plasmids are not sterile. For experiments where strict sterility is required, filtration with

0.22um filter is required.

**RefSeq:** <u>NM 003352.8</u>

 RefSeq Size:
 1527 bp

 RefSeq ORF:
 306 bp

 Locus ID:
 7341

 UniProt ID:
 P63165

 Cytogenetics:
 2q33.1

Domains: UBQ

**Protein Families:** Druggable Genome, Stem cell - Pluripotency, Transcription Factors


**MW:** 11.6 kDa

**Gene Summary:** This gene encodes a protein that is a member of the SUMO (small ubiquitin-like modifier)


protein family. It functions in a manner similar to ubiquitin in that it is bound to target proteins as part of a post-translational modification system. However, unlike ubiquitin which targets proteins for degradation, this protein is involved in a variety of cellular processes, such as nuclear transport, transcriptional regulation, apoptosis, and protein stability. It is not active until the last four amino acids of the carboxy-terminus have been cleaved off. Several pseudogenes have been reported for this gene. Alternate transcriptional splice variants encoding different isoforms have been characterized. [provided by RefSeq, Jul 2008]



# **Product images:**



Circular map for RC200633



Western blot validation of overexpression lysate (Cat# [LY423653]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with [RC221677] using transfection reagent MegaTran 2.0 (Cat# [TT210002]).