

Product datasheet for RC200653L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

NDUFV2 (NM_021074) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: NDUFV2 (NM_021074) Human Tagged ORF Clone Lentiviral Particle

Symbol: NDUFV2

Synonyms: CI-24k; MC1DN7

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_021074

ORF Size: 747 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC200653).

Sequence:

Domains:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 021074.1

 RefSeq Size:
 937 bp

 RefSeq ORF:
 750 bp

 Locus ID:
 4729

 UniProt ID:
 P19404

Cytogenetics: 18p11.22

Protein Families: Druggable Genome

complex1_24kD



NDUFV2 (NM_021074) Human Tagged ORF Clone Lentiviral Particle - RC200653L4V

Protein Pathways: Alzheimer's disease, Huntington's disease, Metabolic pathways, Oxidative phosphorylation,

Parkinson's disease

MW: 27.4 kDa

Gene Summary: The NADH-ubiquinone oxidoreductase complex (complex I) of the mitochondrial respiratory

chain catalyzes the transfer of electrons from NADH to ubiquinone, and consists of at least 43 subunits. The complex is located in the inner mitochondrial membrane. This gene encodes the 24 kDa subunit of complex I, and is involved in electron transfer. Mutations in this gene are implicated in Parkinson's disease, bipolar disorder, schizophrenia, and have been found in one case of early onset hypertrophic cardiomyopathy and encephalopathy. A non-

transcribed pseudogene of this locus is found on chromosome 19. [provided by RefSeq, Oct

2009]