

Product datasheet for RC200663L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Melanoma gp100 (PMEL) (NM_006928) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Melanoma gp100 (PMEL) (NM_006928) Human Tagged ORF Clone Lentiviral Particle

Symbol: Melanoma gp100

Synonyms: D12S53E; gp100; ME20; ME20-M; ME20M; P1; P100; PMEL17; SI; SIL; SILV

Mammalian Cell

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK
ACCN: NM 006928

ORF Size: 1983 bp

ORF Nucleotide

_. _.

Sequence:
OTI Disclaimer:

The ORF insert of this clone is exactly the same as(RC200663).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 006928.3

 RefSeq Size:
 2143 bp

 RefSeq ORF:
 1986 bp

 Locus ID:
 6490

 UniProt ID:
 P40967

Cytogenetics: 12q13.2

Protein Families: Secreted Protein, Transmembrane

MW: 70.1 kDa





Melanoma gp100 (PMEL) (NM_006928) Human Tagged ORF Clone Lentiviral Particle – RC200663L1V

Gene Summary:

This gene encodes a melanocyte-specific type I transmembrane glycoprotein. The encoded protein is enriched in melanosomes, which are the melanin-producing organelles in melanocytes, and plays an essential role in the structural organization of premelanosomes. This protein is involved in generating internal matrix fibers that define the transition from Stage I to Stage II melanosomes. This protein undergoes a complex pattern of prosttranslational processing and modification that is essential to the proper functioning of the protein. A secreted form of this protein that is released by proteolytic ectodomain shedding may be used as a melanoma-specific serum marker. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Jan 2011]