

Product datasheet for **RC200664**

CD137 (TNFRSF9) (NM_001561) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	CD137 (TNFRSF9) (NM_001561) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	CD137
Synonyms:	4-1BB; CD137; CDw137; ILA
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>RC200664 representing NM_001561 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGC**C

ATGGGAACAGCTGTTACAACATAGTAGCCACTCTGTTGCTGGTCTCACTTTGAGAGGACAAGATCAT
TGCAGGATCCTTGTAGTAACTGCCAGCTGGTACATTCTGTGATAATAACAGGAATCAGATTTGCAGTCC
CTGTCTCCAAATAGTTTCTCCAGCGCAGGTGGACAAAGGACCTGTGACATATGCAGGCAGTGTAAAGGT
GTTTTTCAGGACCAGGAAGGAGTGTCTCCACCAGCAATGCAGAGTGTGACTGCACTCCAGGTTTCACT
GCCTGGGGCAGGATGCAGCATGTGTGAACAGGATTGTAACAAGGTCAAGAAGTACAAAAAAGTTG
TAAAGACTGTTGCTTTGGGACATTTAACGATCAGAAACGTGGCATCTGTGACCCCTGGACAACTGTTCT
TTGGATGGAAAGTCTGTGCTTGTGAATGGGACGAAGGAGAGGGACGTGGTCTGTGGACCATCTCCAGCCG
ACCTCTCTCCGGGAGCATCCTCTGTGACCCCGCTGCCCTGCGAGAGAGCCAGGACACTCTCCGCAGAT
CATCTCCTTTCTTTCGCTGACGTCGACTGCGTTGCTCTTCTGCTGTTCTTCTCACGCTCCGTTTC
TCTGTTGTTAAACGGGGCAGAAAGAACTCCTGTATATATTCAAACAACCATTTATGAGACCAGTACAAA
CTACTCAAGAGGAAGATGGCTGTAGTCCGATTTCCAGAAGAAGAAGGAGGATGTGAATG

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)

Protein Sequence: >RC200664 representing NM_001561
 Red=Cloning site Green=Tags(s)

MGNSCYNIVATLLLVLNFERTRSLQDPCSNCPAGTFCDNRRNQICSPCPPNSFSSAGGQRTCDICRQCKG
 VFRTRKECSSTSNAECDCTPGFHCLGAGCSMCEQDCKQGQELTKKGCKDCCFGTNDQKRGICRPWTNCS
 LDGKSVLVNGTKERDVVCGPSPADLSPGASSVTPPAPAREPGHSPQIISFFLALSTALLFLLFLLTRF
 SVVKRGRKLLYIFKQPFMRPVQTTQEEDGCSCRFPEEEEEGGCEL

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites: SgfI-MluI

Cloning Scheme:

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF

ACCN: NM_001561

ORF Size: 765 bp


OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)


OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001561.6
RefSeq Size:	1935 bp
RefSeq ORF:	768 bp
Locus ID:	3604
UniProt ID:	Q07011
Cytogenetics:	1p36.23
Protein Families:	Druggable Genome, Transmembrane
Protein Pathways:	Cytokine-cytokine receptor interaction
MW:	27.9 kDa
Gene Summary:	The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor contributes to the clonal expansion, survival, and development of T cells. It can also induce proliferation in peripheral monocytes, enhance T cell apoptosis induced by TCR/CD3 triggered activation, and regulate CD28 co-stimulation to promote Th1 cell responses. The expression of this receptor is induced by lymphocyte activation. TRAF adaptor proteins have been shown to bind to this receptor and transduce the signals leading to activation of NF-kappaB. [provided by RefSeq, Jul 2008]

Product images:



Circular map for RC200664



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY TNFRSF9 (Cat# RC200664, Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-TNFRSF9 (Cat# [TA813073])(1:500).