

## Product datasheet for RC200699L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## PTPN7 (NM\_002832) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

Product Name: PTPN7 (NM 002832) Human Tagged ORF Clone Lentiviral Particle

Symbol: PTPN7

**Synonyms:** BPTP-4; HEPTP; LC-PTP; LPTP; PTPNI

**Mammalian Cell** 

Selection:

None

**Vector:** pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK
ACCN: NM 002832

ORF Size: 1197 bp

**ORF Nucleotide** 

Sequence:

The ORF insert of this clone is exactly the same as(RC200699).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 002832.2, NP 002823.2

 RefSeq Size:
 3772 bp

 RefSeq ORF:
 1083 bp

 Locus ID:
 5778

 UniProt ID:
 P35236

Cytogenetics: 1q32.1

**Protein Families:** Druggable Genome, Phosphatase

**Protein Pathways:** MAPK signaling pathway





MW:

ORIGENE

45 kDa

**Gene Summary:** 

The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This gene is preferentially expressed in a variety of hematopoietic cells, and is an early response gene in lymphokine stimulated cells. The non-catalytic N-terminus of this PTP can interact with MAP kinases and suppress the MAP kinase activities. This PTP was shown to be involved in the regulation of T cell antigen receptor (TCR) signaling, which was thought to function through dephosphorylating the molecules related to MAP kinase pathway. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Dec 2010]