

Product datasheet for **RC200804**

C3orf10 (BRK1) (NM_018462) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: C3orf10 (BRK1) (NM_018462) Human Tagged ORF Clone
Tag: Myc-DDK
Symbol: C3orf10
Synonyms: C3orf10; hHBrk1; HSPC300; MDS027
Mammalian Cell Selection: Neomycin
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
ORF Nucleotide Sequence: >RC200804 ORF sequence
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGGCGGGACAGGAGGATCCGGTGCAGCGGGAGATTACCAGGACTGGGCTAACCGGGAGTACATTGAGA
TAATCACCAGCAGCATCAAGAAAATCGCAGACTTTCTCAACTCGTTCGATATGCTTGTCTGTTCAAGACT
TGCAACACTAAACGAGAAATTGACAGCCCTTGAACGGAGAATAGAGTACATTGAAGCTCGGGTGACAAA
GGTGAGACTCACC

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC200804 protein sequence
Red=Cloning site Green=Tags(s)
MAGQEDPVQREIHQDWANREYIEIITSSIKKIADFLNSFDMSCRSRLATLNEKLTALERRIEYIEARVTK
GETLT

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mk6429_a12.zip

Restriction Sites: Sgfl-MluI



[View online »](#)

Cloning Scheme:


ACCN: NM_018462

ORF Size: 225 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

- Reconstitution Method:**
1. Centrifuge at 5,000xg for 5min.
 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
 3. Close the tube and incubate for 10 minutes at room temperature.
 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.


Note: Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.

RefSeq: [NM_018462.5](#)


RefSeq Size: 1197 bp

RefSeq ORF:	228 bp
Locus ID:	55845
UniProt ID:	Q8WUW1
Cytogenetics:	3p25.3
Protein Pathways:	Regulation of actin cytoskeleton
MW:	8.7 kDa
Gene Summary:	Involved in regulation of actin and microtubule organization. Part of a WAVE complex that activates the Arp2/3 complex. As component of the WAVE1 complex, required for BDNF-NTRK2 endocytic trafficking and signaling from early endosomes (By similarity). [UniProtKB/Swiss-Prot Function]


Product images:



Circular map for RC200804



Western blot validation of overexpression lysate (Cat# [LY413040]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC200804 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).



Coomassie blue staining of purified BRK1 protein (Cat# [TP300804]). The protein was produced from HEK293T cells transfected with BRK1 cDNA clone (Cat# RC200804) using MegaTran 2.0 (Cat# [TT210002]).