

Product datasheet for **RC200805L3V**

KLF13 (NM_015995) Human Tagged ORF Clone Lentiviral Particle

Product data:

| | |
|---------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Product Type: | Lentiviral Particles |
| Product Name: | KLF13 (NM_015995) Human Tagged ORF Clone Lentiviral Particle |
| Symbol: | KLF13 |
| Synonyms: | BTEB3; FKLF2; NSLP1; RFLAT-1; RFLAT1 |
| Mammalian Cell Selection: | Puromycin |
| Vector: | pLenti-C-Myc-DDK-P2A-Puro (PS100092) |
| Tag: | Myc-DDK |
| ACCN: | NM_015995 |
| ORF Size: | 864 bp |
| ORF Nucleotide Sequence: | The ORF insert of this clone is exactly the same as(RC200805). |
| OTI Disclaimer: | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info |
| OTI Annotation: | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene. |
| RefSeq: | NM_015995.2 |
| RefSeq Size: | 6817 bp |
| RefSeq ORF: | 867 bp |
| Locus ID: | 51621 |
| UniProt ID: | Q9Y2Y9 |
| Cytogenetics: | 15q13.3 |
| Domains: | zf-C2H2 |
| Protein Families: | Druggable Genome, Transcription Factors |


[View online »](#)

MW: 31 kDa

Gene Summary: KLF13 belongs to a family of transcription factors that contain 3 classical zinc finger DNA-binding domains consisting of a zinc atom tetrahedrally coordinated by 2 cysteines and 2 histidines (C2H2 motif). These transcription factors bind to GC-rich sequences and related GT and CACCC boxes (Schohy et al., 2000 [PubMed 11087666]).[supplied by OMIM, Mar 2008]