

Product datasheet for RC200841L1

TTLL12 (NM_015140) Human Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	TTLL12 (NM_015140) Human Tagged Lenti ORF Clone
Tag:	Myc-DDK
Symbol:	TTLL12
Synonyms:	dj526114.2
Mammalian Cell Selection:	None
Vector:	pLenti-C-Myc-DDK (PS100064)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC200841).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF.


ACCN:	NM_015140
ORF Size:	1932 bp




[View online >](#)

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_015140.2
RefSeq Size:	3421 bp
RefSeq ORF:	1935 bp
Locus ID:	23170
UniProt ID:	Q14166
Cytogenetics:	22q13.2
MW:	74.4 kDa
Gene Summary:	Negatively regulates post-translational modifications of tubulin, including detyrosination of the C-terminus and polyglutamylation of glutamate residues (PubMed:20162578, PubMed:23251473). Also, indirectly promotes histone H4 trimethylation at 'Lys-20' (H4K20me3) (PubMed:23251473). Probably by controlling tubulin and/or histone H4 post-translational modifications, plays a role in mitosis and in maintaining chromosome number stability (PubMed:20162578, PubMed:23251473). During RNA virus-mediated infection, acts as a negative regulator of the DDX58/RIG-I pathway by preventing MAVS binding to TBK1 and IKBKE (PubMed:28011935).[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for RC200841L1



Double digestion of RC200841L1 using SgfI and MluI