

Product datasheet for RC200894L3

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

C3IP1 (KLHL12) (NM_021633) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: C3IP1 (KLHL12) (NM_021633) Human Tagged Lenti ORF Clone

Tag: Myc-DDK

Symbol: C3IP1

Synonyms: C3IP1; DKIR **Mammalian Cell** Puromycin

Selection:

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Sgfl-Mlul

E. coli Selection: Chloramphenicol (34 ug/mL)


The ORF insert of this clone is exactly the same as(RC200894). **ORF Nucleotide**

Sequence:

Restriction Sites: Cloning Scheme:

Cloning sites used for ORF Shuttling:





^{*} The last codon before the Stop codon of the ORF.

ACCN: NM_021633

ORF Size: 1704 bp





OTI Disclaimer:

Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>

OTI Annotation:

This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components:

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: <u>NM 021633.2</u>

RefSeq Size: 3338 bp
RefSeq ORF: 1707 bp
Locus ID: 59349
UniProt ID: Q53G59
Cytogenetics: 1q32.1
Domains: BTB, Kelch

MW: 63.3 kDa

Gene Summary:

been identified as an autoantigen in the autoimmune disease Sjogren's syndrome and as a potential biomarker in primary biliary cirrhosis. This protein may act as a substrate adaptor of the Cullin-3 ubiquitin ligase complex to promote substrate-specific ubiquitylation. Ubiquitylation by this complex has been shown to regulate the Wnt signaling pathway as well as COPII vesicle coat size. A pseudogene has been identified on chromosome 22. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Dec 2014]

This gene encodes a member of the KLHL (Kelch-like) family of proteins. This protein has