

Product datasheet for RC200938

Transmembrane protein 93 (EMC6) (NM_031298) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: Transmembrane protein 93 (EMC6) (NM_031298) Human Tagged ORF Clone
Tag: Myc-DDK
Symbol: Transmembrane protein 93
Synonyms: RAB5IFL; TMEM93
Mammalian Cell Selection: Neomycin
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
ORF Nucleotide Sequence: >RC200938 ORF sequence
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCCCGGATCGCC

ATGGCCGCGGTGGTGGCCAAGCGGGAAGGGCCGCCGTTTCATCAGCGAGGCGGCCGTGCGGGGCAACGCCG
 CCGTCTGGATTATTGCCGGACCTCGGTGTGAGCGCTGTCGGGGGCCACGGCCGGCATCTCGGCCCTCAC
 CGGCCTTACGGCTTTCATCTTCTACCTGCTCGCCTCCGTCCTGCTCTCCCTGCTCCTCATTCTCAAGGCG
 GGAAGGAGGTGGAACAAATATTTCAAATCACGGAGACCTCTTTACAGGAGGCCTCATCGGGGCTCT
 TCACCTACGTCTGTTCTGGACGTTCTCTACGGCATGGTGCACGCTAC

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC200938 protein sequence
 Red=Cloning site Green=Tags(s)
 MAAVVAKREGPPFISEAAVRGNAAVLDYCRTSVSALSGATAGILGLTGLYGFIFYLLASVLLSLLLILKA
 GRRWNKYFKSRRPLFTGGLIGLFTYVLFWTFLYGMVHVY

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mk6400_e09.zip



Restriction Sites: SgfI-MluI



[View online »](#)

Cloning Scheme:

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF

ACCN: NM_031298

ORF Size: 330 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

Note: Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.

RefSeq: [NM_031298.4](#)


RefSeq Size: 749 bp

RefSeq ORF: 333 bp


Locus ID: 83460
UniProt ID: [Q9BV81](#)
Cytogenetics: 17p13.2
Protein Families: Transmembrane
MW: 12 kDa

Gene Summary: Part of the endoplasmic reticulum membrane protein complex (EMC) that enables the energy-independent insertion into endoplasmic reticulum membranes of newly synthesized membrane proteins (PubMed:30415835, PubMed:29809151, PubMed:29242231, PubMed:32459176, PubMed:32439656). Preferentially accommodates proteins with transmembrane domains that are weakly hydrophobic or contain destabilizing features such as charged and aromatic residues (PubMed:30415835, PubMed:29809151, PubMed:29242231). Involved in the cotranslational insertion of multi-pass membrane proteins in which stop-transfer membrane-anchor sequences become ER membrane spanning helices (PubMed:30415835, PubMed:29809151). It is also required for the post-translational insertion of tail-anchored/TA proteins in endoplasmic reticulum membranes (PubMed:29809151, PubMed:29242231). By mediating the proper cotranslational insertion of N-terminal transmembrane domains in an N-exo topology, with translocated N-terminus in the lumen of the ER, controls the topology of multi-pass membrane proteins like the G protein-coupled receptors (PubMed:30415835). By regulating the insertion of various proteins in membranes, it is indirectly involved in many cellular processes (Probable).
 [UniProtKB/Swiss-Prot Function]

Product images:



Circular map for RC200938



Western blot validation of overexpression lysate (Cat# [LY423076]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with [RC215548] using transfection reagent MegaTran 2.0 (Cat# [TT210002]).