

Product datasheet for RC201055L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

WDYHV1 (NTAQ1) (NM_018024) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: WDYHV1 (NTAQ1) (NM 018024) Human Tagged ORF Clone Lentiviral Particle

Symbol: NTAQ1

Synonyms: C8orf32; WDYHV1

Mammalian Cell

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK
ACCN: NM 018024

ORF Size: 615 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC201055).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 018024.1</u>

RefSeq Size: 1568 bp
RefSeq ORF: 618 bp
Locus ID: 55093
UniProt ID: Q96HA8
Cytogenetics: 8q24.13

Protein Families: Stem cell - Pluripotency

MW: 23.6 kDa







Gene Summary:

Mediates the side-chain deamidation of N-terminal glutamine residues to glutamate, an important step in N-end rule pathway of protein degradation. Conversion of the resulting N-terminal glutamine to glutamate renders the protein susceptible to arginylation, polyubiquitination and degradation as specified by the N-end rule. Does not act on substrates with internal or C-terminal glutamine and does not act on non-glutamine residues in any position. Does not deaminate acetylated N-terminal glutamine. With the exception of proline, all tested second-position residues on substrate peptides do not greatly influence the activity. In contrast, a proline at position 2, virtually abolishes deamidation of N-terminal glutamine. [UniProtKB/Swiss-Prot Function]