

Product datasheet for RC201075L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

STAT1 (NM_139266) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: STAT1 (NM_139266) Human Tagged ORF Clone Lentiviral Particle

Symbol: STAT

Synonyms: CANDF7; IMD31A; IMD31B; IMD31C; ISGF-3; STAT91

Mammalian Cell

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

Tag:Myc-DDKACCN:NM_139266

ORF Size: 2136 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC201075).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 139266.1</u>

 RefSeq Size:
 2798 bp

 RefSeq ORF:
 2139 bp

 Locus ID:
 6772

 UniProt ID:
 P42224

 Cytogenetics:
 2q32.2

Domains: SH2, STAT

Protein Families: Druggable Genome, Transcription Factors





STAT1 (NM_139266) Human Tagged ORF Clone Lentiviral Particle - RC201075L1V

Protein Pathways: Chemokine signaling pathway, Jak-STAT signaling pathway, Pancreatic cancer, Pathways in

cancer, Toll-like receptor signaling pathway

MW: 83 kDa

Gene Summary: The protein encoded by this gene is a member of the STAT protein family. In response to

cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. The protein encoded by this gene can be activated by various ligands including interferon-alpha, interferon-gamma, EGF, PDGF and IL6. This protein mediates the expression of a variety of genes, which is thought to be important for cell viability in response to different cell stimuli and pathogens. The protein plays an important role in immune responses to viral, fungal and mycobacterial pathogens. Mutations in this gene are associated with Immunodeficiency 31B, 31A, and 31C. [provided by RefSeq,

Jun 2020]