

## Product datasheet for RC201111L2V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Cortactin (CTTN) (NM\_138565) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** Cortactin (CTTN) (NM\_138565) Human Tagged ORF Clone Lentiviral Particle

Symbol: Cortactin

**Synonyms:** EMS1

Mammalian Cell None

Selection:

Vector:

pLenti-C-mGFP (PS100071)

Tag: mGFP

**ACCN:** NM\_138565 **ORF Size:** 1539 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC201111).

•

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 138565.1

 RefSeq Size:
 3208 bp

 RefSeq ORF:
 1542 bp

 Locus ID:
 2017

 UniProt ID:
 Q14247

 Cytogenetics:
 11q13.3

**Protein Families:** Druggable Genome

**Protein Pathways:** Pathogenic Escherichia coli infection, Tight junction





ORIGENE

**MW:** 57.5 kDa

**Gene Summary:** 

This gene is overexpressed in breast cancer and squamous cell carcinomas of the head and neck. The encoded protein is localized in the cytoplasm and in areas of the cell-substratum contacts. This gene has two roles: (1) regulating the interactions between components of adherens-type junctions and (2) organizing the cytoskeleton and cell adhesion structures of epithelia and carcinoma cells. During apoptosis, the encoded protein is degraded in a caspase-dependent manner. The aberrant regulation of this gene contributes to tumor cell invasion and metastasis. Three splice variants that encode different isoforms have been identified for this gene. [provided by RefSeq, May 2010]