

## Product datasheet for RC201113L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Rapsyn (RAPSN) (NM\_032645) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** Rapsyn (RAPSN) (NM\_032645) Human Tagged ORF Clone Lentiviral Particle

Symbol: Rapsyn

Synonyms: CMS4C; CMS11; FADS; FADS2; RAPSYN; RNF205

**Mammalian Cell** 

Selection:

None

**Vector:** pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK
ACCN: NM 032645

ORF Size: 1059 bp

**ORF Nucleotide** 

Sequence:

The ORF insert of this clone is exactly the same as(RC201113).

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 032645.3

 RefSeq Size:
 1494 bp

 RefSeq ORF:
 1062 bp

 Locus ID:
 5913

 UniProt ID:
 Q13702

 Cytogenetics:
 11p11.2

Protein Families: Druggable Genome

MW: 39.9 kDa







## **Gene Summary:**

This gene encodes a member of a family of proteins that are receptor associated proteins of the synapse. The encoded protein contains a conserved cAMP-dependent protein kinase phosphorylation site, and plays a critical role in clustering and anchoring nicotinic acetylcholine receptors at synaptic sites by linking the receptors to the underlying postsynaptic cytoskeleton, possibly by direct association with actin or spectrin. Mutations in this gene may play a role in postsynaptic congenital myasthenic syndromes. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Apr 2011]