

Product datasheet for **RC201117L4V**

TFIIA2 (GTF2A2) (NM_004492) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	TFIIA2 (GTF2A2) (NM_004492) Human Tagged ORF Clone Lentiviral Particle
Symbol:	TFIIA2
Synonyms:	HsT18745; T18745; TF2A2; TFIIA; TFIIA-12; TFIIA-gamma; TFIAS
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_004492
ORF Size:	327 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC201117).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_004492.1
RefSeq Size:	1582 bp
RefSeq ORF:	330 bp
Locus ID:	2958
UniProt ID:	P52657
Cytogenetics:	15q22.2
Domains:	TFIIA_gamma
Protein Families:	Transcription Factors



[View online »](#)

Protein Pathways: Basal transcription factors

MW: 12.5 kDa

Gene Summary: Accurate transcription initiation on TATA-containing class II genes involves the ordered assembly of RNA polymerase II (POLR2A; MIM 180660) and the general initiation factors TFIIA, TFIIB (MIM 189963), TFIID (MIM 313650), TFIIE (MIM 189962), TFIIF (MIM 189968), TFIIG/TFIJ, and TFIIH (MIM 189972). The first step involves recognition of the TATA element by the TATA-binding subunit (TBP; MIM 600075) and may be regulated by TFIIA, a factor that interacts with both TBP and a TBP-associated factor (TAF; MIM 600475) in TFIID. TFIIA has 2 subunits (43 and 12 kD) in yeast and 3 subunits in higher eukaryotes. In HeLa extracts, it consists of a 35-kD alpha subunit and a 19-kD beta subunit encoded by the N- and C-terminal regions of GTF2A1 (MIM 600520), respectively, and a 12-kD gamma subunit encoded by GTF2A2 (DeJong et al., 1995 [PubMed 7724559]).[supplied by OMIM, Mar 2008]