

# **Product datasheet for RC201119**

## DUSP3 (NM\_004090) Human Tagged ORF Clone

### **Product data:**

**Product Type:** Expression Plasmids

Tag: Myc-DDK
Symbol: DUSP3
Synonyms: VHR

Mammalian Cell Neomycin

Selection:

Vector:

pCMV6-Entry (PS100001)

E. coli Selection: Kanamycin (25 ug/mL)

ORF Nucleotide >RC201119 ORF sequence

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ATGTCGGGCTCGTTCGAGCTCTCGGTGCAGGATCTCAACGACCTGCTCTCGGACGGCAGCGGCTGCTACA
GCCTCCCGAGCCAGCCCTGCAACGAGGTCACCCCGCGGATCTACGTGGGCAACGCGTCTGTGGCTCAGGA
CATCCCCAAGCTGCAGAAACTAGGCATCACCCATGTGCTGAACGCGGCTGAGGGCAGGTCCTTCATGCAC
GTCAACACCAATGCCAACTTCTACAAGGACTCCGGCATCACATACCTGGGCATCAAGGCCAACGACACAC
AGGAGTTCAACCTCAGCGCTTACTTTGAAAGGGCTGCCGACTTCATTGACCAGGCTTTGGCTCAAAAGAA
TGGCCGGGTGCTCGTCCACTGCCGGGAAGGTTATAGCCGCTCCCCAACGCTAGTTATCGCCTACCTCATG
ATGCGGCAGAAGATGGACGTCAAGTCTGCCCTGAGCATCGTGAGGCAGAACCGTGAGATCGGCCCCAACG
ATGGCTTCCTGGCCCAGCTCTGCCAGCTCAATGACAGACTAGCCAAGGAGGGGAAGTTGAAACCC

**AGCGGACCG**ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCC

TGGATTACAAGGATGACGACGATAAGGTTTAA

**Protein Sequence:** >RC201119 protein sequence

Red=Cloning site Green=Tags(s)

MSGSFELSVQDLNDLLSDGSGCYSLPSQPCNEVTPRIYVGNASVAQDIPKLQKLGITHVLNAAEGRSFMH VNTNANFYKDSGITYLGIKANDTQEFNLSAYFERAADFIDQALAQKNGRVLVHCREGYSRSPTLVIAYLM

MRQKMDVKSALSIVRQNREIGPNDGFLAQLCQLNDRLAKEGKLKP

**SGPTRTRPL**EQKLISEEDLAANDILDYKDDDDK**V** 

Chromatograms: <a href="https://cdn.origene.com/chromatograms/mk6183">https://cdn.origene.com/chromatograms/mk6183</a> d03.zip

**Restriction Sites:** Sgfl-Rsrll




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



#### **Cloning Scheme:**





<sup>\*</sup> The last codon before the Stop codon of the ORF

**ACCN:** NM\_004090

ORF Size: 555 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**Note:** Plasmids are not sterile. For experiments where strict sterility is required, filtration with

0.22um filter is required.

RefSeq: <u>NM 004090.4</u>

RefSeq Size: 4139 bp
RefSeq ORF: 558 bp
Locus ID: 1845



UniProt ID:P51452Cytogenetics:17q21.31Domains:DSPc

**Protein Families:** Druggable Genome, Phosphatase

**Protein Pathways:** MAPK signaling pathway


MW: 20.5 kDa

**Gene Summary:** The protein encoded by this gene is a member of the dual specificity protein phosphatase

subfamily. These phosphatases inactivate their target kinases by dephosphorylating both the phosphoserine/threonine and phosphotyrosine residues. They negatively regulate members of the mitogen-activated protein (MAP) kinase superfamily (MAPK/ERK, SAPK/JNK, p38), which are associated with cellular proliferation and differentiation. Different members of the family of dual specificity phosphatases show distinct substrate specificities for various MAP kinases, different tissue distribution and subcellular localization, and different modes of inducibility of their expression by extracellular stimuli. This gene maps in a region that contains the BRCA1 locus which confers susceptibility to breast and ovarian cancer. Although DUSP3 is expressed in both breast and ovarian tissues, mutation screening in breast cancer pedigrees and in sporadic tumors was negative, leading to the conclusion that this gene is not BRCA1.


[provided by RefSeq, Jul 2008]


### **Product images:**




Circular map for RC201119










Coomassie blue staining of purified DUSP3 protein (Cat# [TP301119]). The protein was produced from HEK293T cells transfected with DUSP3 cDNA clone (Cat# RC201119) using MegaTran 2.0 (Cat# [TT210002]).



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY DUSP3 (Cat# RC201119, Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-DUSP3 (Cat# [TA812964])(1:500). Positive lysates [LY401320] (100ug) and [LC401320] (20ug) can be purchased separately from OriGene.