

Product datasheet for RC201141L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PRC1 (NM_003981) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: PRC1 (NM_003981) Human Tagged ORF Clone Lentiviral Particle

Symbol: PRC1
Synonyms: ASE1
Mammalian Cell None

Selection:

Vector:

pLenti-C-Myc-DDK (PS100064)

 Tag:
 Myc-DDK

 ACCN:
 NM_003981

 ORF Size:
 1860 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC201141).

OTI Disclaimer: Th

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 003981.2</u>

 RefSeq Size:
 3207 bp

 RefSeq ORF:
 1863 bp

 Locus ID:
 9055

 UniProt ID:
 043663

 Cytogenetics:
 15q26.1

Domains: MAP65_ASE1

MW: 71.7 kDa







Gene Summary:

This gene encodes a protein that is involved in cytokinesis. The protein is present at high levels during the S and G2/M phases of mitosis but its levels drop dramatically when the cell exits mitosis and enters the G1 phase. It is located in the nucleus during interphase, becomes associated with mitotic spindles in a highly dynamic manner during mitosis, and localizes to the cell mid-body during cytokinesis. This protein has been shown to be a substrate of several cyclin-dependent kinases (CDKs). It is necessary for polarizing parallel microtubules and concentrating the factors responsible for contractile ring assembly. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jun 2012]