

Product datasheet for RC201144L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Phosphoribosyl pyrophosphate amidotransferase (PPAT) (NM_002703) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Phosphoribosyl pyrophosphate amidotransferase (PPAT) (NM_002703) Human Tagged ORF

Clone Lentiviral Particle

Symbol: Phosphoribosyl pyrophosphate amidotransferase

Synonyms: ATASE; GPAT; PRAT

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_002703 **ORF Size:** 1551 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC201144).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 002703.3</u>

 RefSeq Size:
 3713 bp

 RefSeq ORF:
 1554 bp

 Locus ID:
 5471

 UniProt ID:
 Q06203

Cytogenetics: 4q12

Domains: GATase_2, Pribosyltran







Protein Families: Druggable Genome, Protease

Protein Pathways: Alanine, aspartate and glutamate metabolism, Metabolic pathways, Purine metabolism

MW: 57.4 kDa

Gene Summary: The protein encoded by this gene is a member of the purine/pyrimidine

phosphoribosyltransferase family. It is a regulatory allosteric enzyme that catalyzes the first step of de novo purine nucleotide biosythetic pathway. This gene and PAICS/AIRC gene, a bifunctional enzyme catalyzing steps six and seven of this pathway, are located in close proximity on chromosome 4, and divergently transcribed from an intergenic region.

[provided by RefSeq, Mar 2011]