

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for RC201168L2V

## Beta Arrestin 2 (ARRB2) (NM\_004313) Human Tagged ORF Clone Lentiviral Particle

## **Product data:**

Product Type:	Lentiviral Particles
Product Name:	Beta Arrestin 2 (ARRB2) (NM_004313) Human Tagged ORF Clone Lentiviral Particle
Symbol:	Beta Arrestin 2
Synonyms:	ARB2; ARR2; BARR2
Mammalian Cell Selection:	None
Vector:	pLenti-C-mGFP (PS100071)
Tag:	mGFP
ACCN:	NM_004313
ORF Size:	1227 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC201168).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 004313.3</u>
RefSeq Size:	1936 bp
RefSeq ORF:	1230 bp
Locus ID:	409
UniProt ID:	<u>P32121</u>
Cytogenetics:	17p13.2
Domains:	arrestin
Protein Families:	Druggable Genome



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Beta Arrestin 2 (ARRB2) (NM_004313) Human Tagged ORF Clone Lentiviral Particle – RC201168L2V	
Protein Pathwa	Chemokine signaling pathway, Endocytosis, MAPK signaling pathway, Olfactory transduc	tion
MW:	46.1 kDa	
Gene Summary	Members of arrestin/beta-arrestin protein family are thought to participate in agonist- mediated desensitization of G-protein-coupled receptors and cause specific dampening cellular responses to stimuli such as hormones, neurotransmitters, or sensory signals. Arrestin beta 2, like arrestin beta 1, was shown to inhibit beta-adrenergic receptor functi vitro. It is expressed at high levels in the central nervous system and may play a role in th regulation of synaptic receptors. Besides the brain, a cDNA for arrestin beta 2 was isolate from thyroid gland, and thus it may also be involved in hormone-specific desensitization TSH receptors. Multiple alternatively spliced transcript variants encoding different isofor have been found for this gene. [provided by RefSeq, Mar 2012]	ion in he ed of

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US