

Product datasheet for RC201170L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TIP49A (RUVBL1) (NM 003707) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: TIP49A (RUVBL1) (NM_003707) Human Tagged ORF Clone Lentiviral Particle

Symbol: TIP49A

Synonyms: ECP-54; ECP54; INO80H; NMP 238; NMP238; PONTIN; Pontin52; RVB1; TIH1; TIP49; TIP49A

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag:Myc-DDKACCN:NM_003707

ORF Size: 1368 bp

ORF Nucleotide Sequence:

The ORF insert of this clone is exactly the same as(RC201170).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 003707.1

 RefSeq Size:
 1785 bp

 RefSeq ORF:
 1371 bp

 Locus ID:
 8607

 UniProt ID:
 Q9Y265

Cytogenetics: 3q21.3 Domains: AAA

Protein Families: Stem cell - Pluripotency, Transcription Factors





TIP49A (RUVBL1) (NM_003707) Human Tagged ORF Clone Lentiviral Particle - RC201170L3V

Protein Pathways: Wnt signaling pathway

MW: 50.2 kDa

Gene Summary: This gene encodes a protein that has both DNA-dependent ATPase and DNA helicase

activities and belongs to the ATPases associated with diverse cellular activities (AAA+) protein family. The encoded protein associates with several multisubunit transcriptional complexes and with protein complexes involved in both ATP-dependent remodeling and histone modification. Alternate splicing results in multiple transcript variants. [provided by RefSeq,

Jan 2016]