

## Product datasheet for **RC201192L3V**

### PPAP2C (PLPP2) (NM\_003712) Human Tagged ORF Clone Lentiviral Particle

#### Product data:

Product Type:	Lentiviral Particles
Product Name:	PPAP2C (PLPP2) (NM_003712) Human Tagged ORF Clone Lentiviral Particle
Symbol:	PPAP2C
Synonyms:	LPP2; PAP-2c; PAP2-g; PPAP2C
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_003712
ORF Size:	864 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC201192).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_003712.2</a>
RefSeq Size:	1320 bp
RefSeq ORF:	867 bp
Locus ID:	8612
UniProt ID:	<a href="#">O43688</a>
Cytogenetics:	19p13.3
Domains:	acidPPc
Protein Families:	Druggable Genome, Stem cell - Pluripotency, Transmembrane



[View online »](#)

<b>Protein Pathways:</b>	Ether lipid metabolism, Fc gamma R-mediated phagocytosis, Glycerolipid metabolism, Glycerophospholipid metabolism, Metabolic pathways, Sphingolipid metabolism
<b>MW:</b>	32.6 kDa
<b>Gene Summary:</b>	<p>The protein encoded by this gene is a member of the phosphatidic acid phosphatase (PAP) family. PAPs convert phosphatidic acid to diacylglycerol, and function in de novo synthesis of glycerolipids as well as in receptor-activated signal transduction mediated by phospholipase D. This protein is similar to phosphatidic acid phosphatase type 2A (PPAP2A) and type 2B (PPAP2B). All three proteins contain 6 transmembrane regions, and a consensus N-glycosylation site. This protein has been shown to possess membrane associated PAP activity. Three alternatively spliced transcript variants encoding distinct isoforms have been reported. [provided by RefSeq, Jul 2008]</p>