

Product datasheet for **RC201200L3V**

Cip4 (TRIP10) (NM_004240) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Cip4 (TRIP10) (NM_004240) Human Tagged ORF Clone Lentiviral Particle
Symbol:	Cip4
Synonyms:	CIP4; HSTP; STOT; STP; TRIP-10
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_004240
ORF Size:	1635 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC201200).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_004240.2
RefSeq Size:	2033 bp
RefSeq ORF:	1638 bp
Locus ID:	9322
UniProt ID:	Q15642
Cytogenetics:	19p13.3
Protein Families:	Druggable Genome
Protein Pathways:	Insulin signaling pathway



[View online »](#)

MW: 62.6 kDa

Gene Summary: Required for translocation of GLUT4 to the plasma membrane in response to insulin signaling (By similarity). Required to coordinate membrane tubulation with reorganization of the actin cytoskeleton during endocytosis. Binds to lipids such as phosphatidylinositol 4,5-bisphosphate and phosphatidylserine and promotes membrane invagination and the formation of tubules. Also promotes CDC42-induced actin polymerization by recruiting WASL/N-WASP which in turn activates the Arp2/3 complex. Actin polymerization may promote the fission of membrane tubules to form endocytic vesicles. Required for the formation of podosomes, actin-rich adhesion structures specific to monocyte-derived cells. May be required for the lysosomal retention of FASLG/FASL.[UniProtKB/Swiss-Prot Function]