

## Product datasheet for RC201201L2V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## PSMD11 (NM\_002815) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** PSMD11 (NM\_002815) Human Tagged ORF Clone Lentiviral Particle

Symbol: PSMD11

Synonyms: p44.5; Rpn6; S9

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-mGFP (PS100071)

Tag: mGFP

**ACCN:** NM\_002815 **ORF Size:** 1266 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC201201).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 002815.2

 RefSeq Size:
 1598 bp

 RefSeq ORF:
 1269 bp

 Locus ID:
 5717

 UniProt ID:
 000231

 Cytogenetics:
 17q11.2

Domains: PCI

**Protein Families:** Stem cell - Pluripotency







**Protein Pathways:** Proteasome

MW: 47.3 kDa

**Gene Summary:** The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure

composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. This gene encodes a member of the proteasome subunit S9 family that functions as a non-ATPase subunit of the 19S regulator and is phosphorylated by AMP-activated protein kinase.

subunit of the 19S regulator and is phosphorylated by AMP-activated protein kinase.

Alternatively spliced transcript variants have been observed for this gene. [provided by

RefSeq, Jul 2012]