

Product datasheet for RC201224L3

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

MCK10 (DDR1) (NM_001954) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: MCK10 (DDR1) (NM_001954) Human Tagged Lenti ORF Clone

Tag: Myc-DDK Symbol: MCK10

Synonyms: CAK; CD167; DDR; EDDR1; HGK2; MCK10; NEP; NTRK4; PTK3; PTK3A; RTK6; TRKE

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this clor

Sequence:

The ORF insert of this clone is exactly the same as(RC201224).

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF.

ACCN: NM_001954

ORF Size: 2628 bp





MCK10 (DDR1) (NM_001954) Human Tagged Lenti ORF Clone - RC201224L3

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001954.4</u>, <u>NP 001945.3</u>

RefSeq Size:3840 bpRefSeq ORF:2631 bp

Locus ID: 780

UniProt ID: Q08345
Cytogenetics: 6p21.33

Protein Families: Druggable Genome, Protein Kinase, Transmembrane

MW: 97 kDa


Gene Summary: Receptor tyrosine kinases play a key role in the communication of cells with their

microenvironment. These kinases are involved in the regulation of cell growth, differentiation and metabolism. The protein encoded by this gene belongs to a subfamily of tyrosine kinase receptors with homology to Dictyostelium discoideum protein discoidin I in their extracellular domain, and that are activated by various types of collagen. Expression of this protein is restricted to epithelial cells, particularly in the kidney, lung, gastrointestinal tract, and brain. In addition, it has been shown to be significantly overexpressed in several human tumors. Alternatively spliced transcript variants encoding different isoforms have been described for

this gene. [provided by RefSeq, Feb 2011]



Product images:



Double digestion of RC201224L3 using Sgfl and Mlul $\,$