

Product datasheet for RC201238L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

POLD1 (NM_002691) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: POLD1 (NM_002691) Human Tagged ORF Clone Lentiviral Particle

Symbol: POLD1

Synonyms: CDC2; CRCS10; MDPL; POLD

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_002691 **ORF Size:** 3321 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC201238).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 002691.1

 RefSeq Size:
 3464 bp

 RefSeq ORF:
 3324 bp

 Locus ID:
 5424

 UniProt ID:
 P28340

 Cytogenetics:
 19q13.33

Domains: DNA_pol_B_exo, DNA_pol_B, POLBc

Protein Families: Druggable Genome, Stem cell - Pluripotency





POLD1 (NM_002691) Human Tagged ORF Clone Lentiviral Particle - RC201238L4V

Protein Pathways: Base excision repair, DNA replication, Homologous recombination, Metabolic pathways,

Mismatch repair, Nucleotide excision repair, Purine metabolism, Pyrimidine metabolism

MW: 123.6 kDa

Gene Summary: This gene encodes the 125-kDa catalytic subunit of DNA polymerase delta. DNA polymerase

delta possesses both polymerase and 3' to 5' exonuclease activity and plays a critical role in DNA replication and repair. Alternatively spliced transcript variants have been observed for this gene, and a pseudogene of this gene is located on the long arm of chromosome 6.

[provided by RefSeq, Mar 2012]