

## Product datasheet for RC201251L2V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## PSMC5 (NM\_002805) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

Product Name: PSMC5 (NM 002805) Human Tagged ORF Clone Lentiviral Particle

Symbol: PSMC5

Synonyms: p45; p45/SUG; RPT6; S8; SUG-1; SUG1; TBP10; TRIP1

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM\_002805

ORF Size: 1218 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC201251).

Sequence:
OTI Disclaimer:

**Domains:** 

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 002805.4

 RefSeq Size:
 1372 bp

 RefSeq ORF:
 1221 bp

 Locus ID:
 5705

 UniProt ID:
 P62195

 Cytogenetics:
 17q23.3

**Protein Families:** Druggable Genome

AAA, AAA







**Protein Pathways:** Proteasome

MW: 45.6 kDa

**Gene Summary:** The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure

composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes one of the ATPase subunits, a member of the triple-A family of ATPases which have a chaperone-like activity. In addition to participation in proteasome functions, this subunit may participate in transcriptional regulation since it has been shown to interact with the thyroid hormone receptor and retinoid X receptor-alpha. Two transcript variants encoding different isoforms have been found for this gene. [provided]

by RefSeg, Nov 2010]