

Product datasheet for RC201266L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

POLR2E (NM 002695) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: POLR2E (NM_002695) Human Tagged ORF Clone Lentiviral Particle

Symbol:

hRPB25; hsRPB5; RPABC1; RPB5; XAP4 Synonyms:

Mammalian Cell

Selection:

ACCN:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Myc-DDK Tag: NM 002695

ORF Size: 630 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC201266).

Sequence: OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 002695.2

RefSeq Size: 2866 bp RefSeq ORF: 633 bp Locus ID: 5434 **UniProt ID:** P19388 Cytogenetics: 19p13.3

Domains: RNA_pol_Rpb5_C, RNA_pol_Rpb5_N

Protein Families: Transcription Factors





POLR2E (NM_002695) Human Tagged ORF Clone Lentiviral Particle - RC201266L3V

Protein Pathways: Huntington's disease, Metabolic pathways, Purine metabolism, Pyrimidine metabolism, RNA

polymerase

MW: 24.6 kDa

Gene Summary: This gene encodes the fifth largest subunit of RNA polymerase II, the polymerase responsible

for synthesizing messenger RNA in eukaryotes. This subunit is shared by the other two DNA-directed RNA polymerases and is present in two-fold molar excess over the other polymerase subunits. An interaction between this subunit and a hepatitis virus transactivating protein has been demonstrated, suggesting that interaction between transcriptional activators and the polymerase can occur through this subunit. A pseudogene is located on chromosome 11. Three transcript variants encoding two different isoforms have been found for this gene.

[provided by RefSeq, Oct 2015]