

Product datasheet for RC201272L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

BAG1 (NM_004323) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: BAG1 (NM 004323) Human Tagged ORF Clone Lentiviral Particle

Symbol: BAG

Synonyms: BAG-1; HAP; RAP46

Mammalian Cell

Selection:

Puromycin

Vector:

pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_004323 **ORF Size:** 1035 bp

ORF Nucleotide

- - - - 1

Sequence:

The ORF insert of this clone is exactly the same as(RC201272).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 004323.4

RefSeq Size:3885 bpRefSeq ORF:1038 bp

Locus ID: 573

 UniProt ID:
 Q99933

 Cytogenetics:
 9p13.3

Domains: UBQ, BAG

Protein Families: Druggable Genome







MW: 38.9 kDa

Gene Summary: The oncogene BCL2 is a membrane protein that blocks a step in a pathway leading to

apoptosis or programmed cell death. The protein encoded by this gene binds to BCL2 and is referred to as BCL2-associated athanogene. It enhances the anti-apoptotic effects of BCL2 and represents a link between growth factor receptors and anti-apoptotic mechanisms. Multiple protein isoforms are encoded by this mRNA through the use of a non-AUG (CUG) initiation codon, and three alternative downstream AUG initiation codons. A related pseudogene has been defined on chromosome X. [provided by RefSeq, Feb 2010]