

Product datasheet for RC201345L1

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

DOPA Decarboxylase (DDC) (NM_000790) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: DOPA Decarboxylase (DDC) (NM_000790) Human Tagged Lenti ORF Clone

Tag: Myc-DDK

Symbol: DOPA Decarboxylase

Synonyms: AADC

Mammalian Cell None

Selection:

Vector:pLenti-C-Myc-DDK (PS100064)E. coli Selection:Chloramphenicol (34 ug/mL)


ORF Nucleotide The ORF insert of this clone is exactly the same as(RC201345).

Sequence:

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF.

ACCN: NM_000790

ORF Size: 1440 bp





DOPA Decarboxylase (DDC) (NM_000790) Human Tagged Lenti ORF Clone - RC201345L1

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 000790.3</u>, <u>NP 000781.1</u>

 RefSeq Size:
 1975 bp

 RefSeq ORF:
 1443 bp

 Locus ID:
 1644

 UniProt ID:
 P20711

Cytogenetics: 7p12.2-p12.1

Protein Families: Druggable Genome

Protein Pathways: Histidine metabolism, Metabolic pathways, Phenylalanine metabolism, Tryptophan

metabolism, Tyrosine metabolism

MW: 53.9 kDa

Gene Summary: The encoded protein catalyzes the decarboxylation of L-3,4-dihydroxyphenylalanine (DOPA)


to dopamine, L-5-hydroxytryptophan to serotonin and L-tryptophan to tryptamine. Defects in this gene are the cause of aromatic L-amino-acid decarboxylase deficiency (AADCD). AADCD deficiency is an inborn error in neurotransmitter metabolism that leads to combined

encoding different isoforms have been identified for this gene. [provided by RefSeq, Jun 2011]


serotonin and catecholamine deficiency. Multiple alternatively spliced transcript variants



Product images:



Circular map for RC201345L1



Double digestion of RC201345L1 using Sgfl and Mlul