

Product datasheet for RC201348L3

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

EIF4EBP1 (NM_004095) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: EIF4EBP1 (NM_004095) Human Tagged Lenti ORF Clone

Tag: Myc-DDK Symbol: EIF4EBP1

Synonyms: 4E-BP1; 4EBP1; BP-1; PHAS-I

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this clone is exactly the same as(RC201348).

Sequence:

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





st The last codon before the Stop codon of the ORF.

ACCN: NM_004095

ORF Size: 354 bp





OTI Disclaimer:

Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at customercom or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>

OTI Annotation:

This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components:

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: <u>NM 004095.2</u>

RefSeq Size: 895 bp
RefSeq ORF: 357 bp
Locus ID: 1978
UniProt ID: Q13541
Cytogenetics: 8p11.23

Protein Pathways: Acute myeloid leukemia, ErbB signaling pathway, Insulin signaling pathway, mTOR signaling

pathway

MW: 12.4 kDa

Gene Summary: This gene encodes one member of a family of translation repressor proteins. The protein

directly interacts with eukaryotic translation initiation factor 4E (eIF4E), which is a limiting component of the multisubunit complex that recruits 40S ribosomal subunits to the 5' end of mRNAs. Interaction of this protein with eIF4E inhibits complex assembly and represses translation. This protein is phosphorylated in response to various signals including UV irradiation and insulin signaling, resulting in its dissociation from eIF4E and activation of

mRNA translation. [provided by RefSeq, Jul 2008]