

## Product datasheet for RC201377L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## AP2M1 (NM\_001025205) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

Product Name: AP2M1 (NM\_001025205) Human Tagged ORF Clone Lentiviral Particle

Symbol: AP2M1

Synonyms: AP50; CLAPM1; MRD60; mu2

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

**ACCN:** NM\_001025205

ORF Size: 1305 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC201377).

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** NM 001025205.1, NP 001020376.1

RefSeq Size: 1952 bp
RefSeq ORF: 1302 bp
Locus ID: 1173
UniProt ID: Q96CW1

Cytogenetics: 3q27.1

**Protein Families:** Druggable Genome

**Protein Pathways:** Endocytosis, Huntington's disease





ORÏGENE

**MW:** 49.2 kDa

**Gene Summary:** This gene encodes a subunit of the heterotetrameric coat assembly protein complex 2 (AP2),

which belongs to the adaptor complexes medium subunits family. The encoded protein is required for the activity of a vacuolar ATPase, which is responsible for proton pumping occurring in the acidification of endosomes and lysosomes. The encoded protein may also play an important role in regulating the intracellular trafficking and function of CTLA-4 protein. Three transcript variants encoding different isoforms have been found for this gene.

[provided by RefSeq, Jul 2015]