


Product datasheet for RC201393L4

TCEB2 (ELOB) (NM_007108) Human Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	TCEB2 (ELOB) (NM_007108) Human Tagged Lenti ORF Clone
Tag:	mGFP
Symbol:	ELOB
Synonyms:	SIII; TCEB2
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC201393).
Restriction Sites:	Sgfl-Mlul
Cloning Scheme:	
	Cloning sites used for ORF Shuttling:
	Sgf1 ORF Miu I GCG ATC GC/C ATG // NNN ACG CGT



ACCN: ORF Size: NM_007108 354 bp

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More infoOTI Annotation:This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.Components:The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).Reconstitution Method:1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.Note:Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.Refseq:NM 007108.2Refseq ORF:357 bpLocus ID:Q15370Cytogenetics:16p13.3Domains:UBQProtein Families:Druggable Genome, Transcription FactorsProtein Families:Pathways in cancer, Renal cell carcinoma, Ubiquitin mediated proteolysisMW:13.1 kDa	CRIGENE TCEB2 (ELOB) (NM_007108) Human Tagged Lenti ORF Clone – RC201393L4	
varies depending on the nature of the gene.Components:The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).Reconstitution Method:1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.Note:Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.RefSeq:NM 007108.2RefSeq Size:1009 bpRefSeq ORF:357 bpLocus ID:015370Cytogenetics:16p13.3Domains:UBQProtein Families:Druggable Genome, Transcription FactorsProtein Families:Pruggable Genome, Transcription FactorsW:13.1 kDaGene Summary:This gene encodes the protein elongin B, which is a subunit of the transcription factor B (SIII) complex. The SIII complex is composed of elongins A/A2, B and C, It activates elongation by RNA polymerase II by suppressing transient pausing of the polymerase at many sites within transcription units. Elongin A functions as the transcription senced of the SIII complex. whereas elongins B and C are regulatory subunits. Elongins B and C. The von Hippel-Lindau tumor suppressor protein binds to elongins B and C. The von Hippel-Lindau tumor suppressor protein binds to elongins B and C. The <b< td=""><td>OTI Disclaimer:</td><td>reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing</td></b<>	OTI Disclaimer:	reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing
containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).Reconstitution Method:1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of 	OTI Annotation:	
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.Note:Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.RefSeq:NM 007108.2RefSeq Size:1009 bpRefSeq ORF:357 bpLocus ID:015370Cytogenetics:16p13.3Domains:UBQProtein Pathways:Pathways in cancer, Renal cell carcinoma, Ubiquitin mediated proteolysisMW:13.1 kDaGene Summary:This gene encodes the protein elongin B, which is a subunit of the transcription factor B (SIII) complex, the SIII complex is composed of elongins A/A2, B and C. It activates elongation by RNA polymerase II by suppressing transient pausing of the polymerase at many sites within transcription units. Elongin A functions as the transcriptionally active component of the SIII complex, whereas elongins B and C are regulatory subunits. Elongin A 2 is specifically expressed in the testis, and capable of forming a stable complex with elongins B and C. The von Hippel-Lindu tumor suppressor protein binds to elongins B and C, and thereby inhibits transcription elongation. Two alternatively spliced transcript variants encoding different isoforms have been described for this gene. Pseudogenes have been identified on	Components:	
0.22um filter is required.RefSeq:NM 007108.2RefSeq Size:1009 bpRefSeq ORF:357 bpLocus ID:6923UniProt ID:015370Cytogenetics:16p13.3Domains:UBQProtein Families:Druggable Genome, Transcription FactorsProtein Pathways:Pathways in cancer, Renal cell carcinoma, Ubiquitin mediated proteolysisMW:13.1 kDaGene Summary:This gene encodes the protein elongin B, which is a subunit of the transcription factor B (SIII) complex. The SIII complex is composed of elongins A/A2, B and C. It activates elongation by RNA polymerase II by suppressing transient pausing of the polymerase at many sites within transcription units. Elongin A functions as the transcriptionally active component of the SIII complex, whereas elongins B and C are regulatory subunits. Elongin A2 is specifically expressed in the testis, and capable of forming a stable complex with elongins B and C. The von Hippel-Lindau tumor suppressor protein binds to elongins B and C, and thereeby inhibits transcription elongation. Two alternatively spliced transcript variants encoding different isoforms have been described for this gene. Pseudogenes have been identified on	Reconstitution Met	 Carefully open the tube and add 100ul of sterile water to dissolve the DNA. Close the tube and incubate for 10 minutes at room temperature. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of
RefSeq Size:1009 bpRefSeq ORF:357 bpLocus ID:6923UniProt ID:015370Cytogenetics:16p13.3Domains:UBQProtein Families:Druggable Genome, Transcription FactorsProtein Pathways:Pathways in cancer, Renal cell carcinoma, Ubiquitin mediated proteolysisMW:13.1 kDaGene Summary:This gene encodes the protein elongin B, which is a subunit of the transcription pathway is transcription units. Elongin A functionally active component of the SIII complex. The SIII complex is composed of elongins A/A2, B and C. It activates elongation by RNA polymerase II by suppressing transient pausing of the polymerase at many sites within transcription units. Elongin A functions as the transcriptionally active component of the SIII complex, whereas elongins B and C are regulatory subunits. Elongin A2 is specifically expressed in the testis, and capable of forming a stable complex with elongins B and C. The yon Hippel-Lindau tumor suppressor protein binds to elongins B and C, and thereby inhibits transcription elongation. Two alternatively spliced transcript variants encoding different isoforms have been described for this gene. Pseudogenes have been identified on	Note:	
RefSeq ORF:357 bpLocus ID:6923UniProt ID:Q15370Cytogenetics:16p13.3Domains:UBQProtein Families:Druggable Genome, Transcription FactorsProtein Pathways:Pathways in cancer, Renal cell carcinoma, Ubiquitin mediated proteolysisMW:13.1 kDaGene Summary:This gene encodes the protein elongin B, which is a subunit of the transcription factor B (SIII) complex. The SIII complex is composed of elongins A/A2, B and C. It activates elongation by RNA polymerase II by suppressing transient pausing of the polymerase at many sites within transcription units. Elongin A functions as the transcriptionally active component of the SIII complex, whereas elongins B and C are regulatory subunits. Elongin A2 is specifically expressed in the testis, and capable of forming a stable complex with elongins B and C. The von Hippel-Lindau tumor suppressor protein binds to elongins B and C, and thereby inhibits transcription elongation. Two alternatively spliced transcript variants encoding different isoforms have been described for this gene. Pseudogenes have been identified on	RefSeq:	<u>NM 007108.2</u>
Locus ID:6923UniProt ID:Q15370Cytogenetics:16p13.3Domains:UBQProtein Families:Druggable Genome, Transcription FactorsProtein Pathways:Pathways in cancer, Renal cell carcinoma, Ubiquitin mediated proteolysisMW:13.1 kDaGene Summary:This gene encodes the protein elongin B, which is a subunit of the transcription factor B (SIII) complex. The SIII complex is composed of elongins A/A2, B and C. It activates elongation by RNA polymerase II by suppressing transient pausing of the polymerase at many sites within transcription units. Elongin A functions as the transcriptionally active component of the SIII complex, whereas elongins B and C are regulatory subunits. Elongin A2 is specifically expressed in the testis, and capable of forming a stable complex with elongins B and C. The von Hippel-Lindau tumor suppressor protein binds to elongins B and C, and thereby inhibits transcription elongation. Two alternatively spliced transcript variants encoding different isoforms have been described for this gene. Pseudogenes have been identified on	RefSeq Size:	1009 bp
UniProt ID:Q15370Cytogenetics:16p13.3Domains:UBQProtein Families:Druggable Genome, Transcription FactorsProtein Pathways:Pathways in cancer, Renal cell carcinoma, Ubiquitin mediated proteolysisMW:13.1 kDaGene Summary:This gene encodes the protein elongin B, which is a subunit of the transcription factor B (SIII) complex. The SIII complex is composed of elongins A/A2, B and C. It activates elongation by RNA polymerase II by suppressing transient pausing of the polymerase at many sites within transcription units. Elongin A functions as the transcriptionally active component of the SIII complex, whereas elongins B and C are regulatory subunits. Elongin A2 is specifically expressed in the testis, and capable of forming a stable complex with elongins B and C. The von Hippel-Lindau tumor suppressor protein binds to elongins B and C, and thereby inhibits transcription elongation. Two alternatively spliced transcript variants encoding different isoforms have been described for this gene. Pseudogenes have been identified on	RefSeq ORF:	357 bp
Cytogenetics:16p13.3Domains:UBQProtein Families:Druggable Genome, Transcription FactorsProtein Pathways:Pathways in cancer, Renal cell carcinoma, Ubiquitin mediated proteolysisMW:13.1 kDaGene Summary:This gene encodes the protein elongin B, which is a subunit of the transcription factor B (SIII) complex. The SIII complex is composed of elongins A/A2, B and C. It activates elongation by RNA polymerase II by suppressing transient pausing of the polymerase at many sites within transcription units. Elongin A functions as the transcriptionally active component of the SIII complex, whereas elongins B and C are regulatory subunits. Elongin A2 is specifically expressed in the testis, and capable of forming a stable complex with elongins B and C. The von Hippel-Lindau tumor suppressor protein binds to elongins B and C, and thereby inhibits transcription elongation. Two alternatively spliced transcript variants encoding different isoforms have been described for this gene. Pseudogenes have been identified on	Locus ID:	6923
Domains:UBQProtein Families:Druggable Genome, Transcription FactorsProtein Pathways:Pathways in cancer, Renal cell carcinoma, Ubiquitin mediated proteolysisMW:13.1 kDaGene Summary:This gene encodes the protein elongin B, which is a subunit of the transcription factor B (SIII) complex. The SIII complex is composed of elongins A/A2, B and C. It activates elongation by RNA polymerase II by suppressing transient pausing of the polymerase at many sites within transcription units. Elongin A functions as the transcriptionally active component of the SIII complex, whereas elongins B and C are regulatory subunits. Elongin A2 is specifically expressed in the testis, and capable of forming a stable complex with elongins B and C. The von Hippel-Lindau tumor suppressor protein binds to elongins B and C, and thereby inhibits transcription elongation. Two alternatively spliced transcript variants encoding different isoforms have been described for this gene. Pseudogenes have been identified on	UniProt ID:	<u>Q15370</u>
Protein Families:Druggable Genome, Transcription FactorsProtein Pathways:Pathways in cancer, Renal cell carcinoma, Ubiquitin mediated proteolysisMW:13.1 kDaGene Summary:This gene encodes the protein elongin B, which is a subunit of the transcription factor B (SIII) complex. The SIII complex is composed of elongins A/A2, B and C. It activates elongation by RNA polymerase II by suppressing transient pausing of the polymerase at many sites within transcription units. Elongin A functions as the transcriptionally active component of the SIII complex, whereas elongins B and C are regulatory subunits. Elongin A2 is specifically expressed in the testis, and capable of forming a stable complex with elongins B and C. The von Hippel-Lindau tumor suppressor protein binds to elongins B and C, and thereby inhibits transcription elongation. Two alternatively spliced transcript variants encoding different isoforms have been described for this gene. Pseudogenes have been identified on	Cytogenetics:	16p13.3
Protein Pathways:Pathways in cancer, Renal cell carcinoma, Ubiquitin mediated proteolysisMW:13.1 kDaGene Summary:This gene encodes the protein elongin B, which is a subunit of the transcription factor B (SIII) complex. The SIII complex is composed of elongins A/A2, B and C. It activates elongation by RNA polymerase II by suppressing transient pausing of the polymerase at many sites within transcription units. Elongin A functions as the transcriptionally active component of the SIII complex, whereas elongins B and C are regulatory subunits. Elongin A2 is specifically expressed in the testis, and capable of forming a stable complex with elongins B and C. The von Hippel-Lindau tumor suppressor protein binds to elongins B and C, and thereby inhibits transcription elongation. Two alternatively spliced transcript variants encoding different isoforms have been described for this gene. Pseudogenes have been identified on	Domains:	UBQ
MW: 13.1 kDa Gene Summary: This gene encodes the protein elongin B, which is a subunit of the transcription factor B (SIII) complex. The SIII complex is composed of elongins A/A2, B and C. It activates elongation by RNA polymerase II by suppressing transient pausing of the polymerase at many sites within transcription units. Elongin A functions as the transcriptionally active component of the SIII complex, whereas elongins B and C are regulatory subunits. Elongin A2 is specifically expressed in the testis, and capable of forming a stable complex with elongins B and C. The von Hippel-Lindau tumor suppressor protein binds to elongins B and C, and thereby inhibits transcription elongation. Two alternatively spliced transcript variants encoding different isoforms have been described for this gene. Pseudogenes have been identified on	Protein Families:	Druggable Genome, Transcription Factors
Gene Summary: This gene encodes the protein elongin B, which is a subunit of the transcription factor B (SIII) complex. The SIII complex is composed of elongins A/A2, B and C. It activates elongation by RNA polymerase II by suppressing transient pausing of the polymerase at many sites within transcription units. Elongin A functions as the transcriptionally active component of the SIII complex, whereas elongins B and C are regulatory subunits. Elongin A2 is specifically expressed in the testis, and capable of forming a stable complex with elongins B and C. The von Hippel-Lindau tumor suppressor protein binds to elongins B and C, and thereby inhibits transcription elongation. Two alternatively spliced transcript variants encoding different isoforms have been described for this gene. Pseudogenes have been identified on	Protein Pathways:	Pathways in cancer, Renal cell carcinoma, Ubiquitin mediated proteolysis
complex. The SIII complex is composed of elongins A/A2, B and C. It activates elongation by RNA polymerase II by suppressing transient pausing of the polymerase at many sites within transcription units. Elongin A functions as the transcriptionally active component of the SIII complex, whereas elongins B and C are regulatory subunits. Elongin A2 is specifically expressed in the testis, and capable of forming a stable complex with elongins B and C. The von Hippel-Lindau tumor suppressor protein binds to elongins B and C, and thereby inhibits transcription elongation. Two alternatively spliced transcript variants encoding different isoforms have been described for this gene. Pseudogenes have been identified on	MW:	13.1 kDa
	Gene Summary:	complex. The SIII complex is composed of elongins A/A2, B and C. It activates elongation by RNA polymerase II by suppressing transient pausing of the polymerase at many sites within transcription units. Elongin A functions as the transcriptionally active component of the SIII complex, whereas elongins B and C are regulatory subunits. Elongin A2 is specifically expressed in the testis, and capable of forming a stable complex with elongins B and C. The von Hippel-Lindau tumor suppressor protein binds to elongins B and C, and thereby inhibits transcription elongation. Two alternatively spliced transcript variants encoding different isoforms have been described for this gene. Pseudogenes have been identified on

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Product images:



Circular map for RC201393L4

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US