

Product datasheet for RC201393L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TCEB2 (ELOB) (NM_007108) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: TCEB2 (ELOB) (NM_007108) Human Tagged ORF Clone Lentiviral Particle

Symbol: ELOB

Synonyms: SIII; TCEB2

Mammalian Cell

Selection:

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Puromycin

Tag: mGFP

ACCN: NM_007108

ORF Size: 354 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC201393).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 007108.2</u>

 RefSeq Size:
 1009 bp

 RefSeq ORF:
 357 bp

 Locus ID:
 6923

 UniProt ID:
 Q15370

 Cytogenetics:
 16p13.3

 Domains:
 UBQ

Protein Families: Druggable Genome, Transcription Factors





TCEB2 (ELOB) (NM_007108) Human Tagged ORF Clone Lentiviral Particle - RC201393L4V

Protein Pathways: Pathways in cancer, Renal cell carcinoma, Ubiquitin mediated proteolysis

MW: 13.1 kDa

Gene Summary: This gene encodes the protein elongin B, which is a subunit of the transcription factor B (SIII)

complex. The SIII complex is composed of elongins A/A2, B and C. It activates elongation by RNA polymerase II by suppressing transient pausing of the polymerase at many sites within transcription units. Elongin A functions as the transcriptionally active component of the SIII complex, whereas elongins B and C are regulatory subunits. Elongin A2 is specifically

expressed in the testis, and capable of forming a stable complex with elongins B and C. The von Hippel-Lindau tumor suppressor protein binds to elongins B and C, and thereby inhibits transcription elongation. Two alternatively spliced transcript variants encoding different

isoforms have been described for this gene. Pseudogenes have been identified on

chromosomes 11 and 13. [provided by RefSeq, Aug 2008]