

Product datasheet for **RC201419**

EXOSC7 (NM_015004) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	EXOSC7 (NM_015004) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	EXOSC7
Synonyms:	EAP1; hRrp42p; p8; RRP42; Rrp42p
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>RC201419 ORF sequence Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGC**C

ATGGCGTCCGTGACGCTGAGCGAGGCGGAGAAGGTGTACATCGTGCATGGCGTCCAGGAAGACCTCCGTG
TGGATGGCCGTGGCTGTGAGGACTACCGATGTGTGCAAGTGAAACTGATGTGGTGTCCAACACTAGTGG
GTCCGCCAGGGTCAAGCTGGGTACACAGACATCTTGGTGGGAGTGAAAGCAGAAATGGGGACGCCGAAG
CTGGAGAAACCAATGAAGGCTACTTGGAGTTCTTTGTTGACTGTTGAGCCAGTGTACCCCTGAATTTG
AAGGTAGAGGAGGTGATGACCTTGGCACCAGATCGCTAACACCCTCTATCGGATATTTAAACAATAAAAG
CAGTGTGCACTTAAAGACCCTCTGCATTAGTCCCTCGGGAGCACTGCTGGTTCTCTATGTGGATGTGCTG
CTTCTGGAATGTGGTGGAAATTTGTTTGTGCCATTTCCATTGCTGTAAGGCTGCTCTCTCAATACAA
GGATACCAAGGGTTCGAGTTTTGGAGGATGAAGAGGGGTGCAAGGACATTGAATTGTCAGATGACCCCTTA
TGACTGCATACGACTAAGTGTGGAGAAATGTCCCCTGCATTGCTCACTCTGTGCAAGATTGGCTATCGGCAT
GTGGTGGATGCTACTCTCAGGAGGAGGCTGCTCGTGGCCAGCTTGTGGTGTGCGTGACCAGCAAGG
GAGTTGTGACGTGCATGAGGAAAGTGGGAAGGGCAGCCTGGACCCAGAGAGCATCTTCGAGATGATGGA
GACTGGCAAGCGTGTGGCAAGGTAAGTGCATGCCTCCTTCAGAGTGTCTGCACAAGGAAGAAAGCCTG
GGCCCAAGAGACAGAAAGTTGGATTCTGGGA

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)

Protein Sequence: >RC201419 protein sequence
Red=Cloning site Green=Tags(s)

MASVTLSEAEKVYIVHGVQEDLRVDGRGCEYRCVEVETDVVSNTSGSARVKLGHTDILVGVAEMGTPK
 LEKPNEGYLEFFVDCSASATPEFEGRGGDDLGTETIANTLYRIFNNKSSVDLKTLCISPREHCWVLYVDVL
 LLECGNLFDAISIAVKAALFNTRIPRVRVLEDEEGSKDIELSDDPYDCIRLSVENVPCIVTLCKIGYRH
 VVDATLQEEACSLASLLVSVTSKGVVTCMRKVGKGLDPESIFEMMETGKRVGKVLHASLQSVLHKEESL
 GPKRQKVGFLG

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mk6053_a12.zip

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_015004

ORF Size: 873 bp


OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.


Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_015004.2 , NP_055819.1
RefSeq Size:	1059 bp
RefSeq ORF:	876 bp
Locus ID:	23016
UniProt ID:	Q15024
Cytogenetics:	3p21.31
Domains:	RNase_PH_C
Protein Families:	Stem cell - Pluripotency
Protein Pathways:	RNA degradation
MW:	31.8 kDa
Gene Summary:	<p>Non-catalytic component of the RNA exosome complex which has 3'->5' exoribonuclease activity and participates in a multitude of cellular RNA processing and degradation events. In the nucleus, the RNA exosome complex is involved in proper maturation of stable RNA species such as rRNA, snRNA and snoRNA, in the elimination of RNA processing by-products and non-coding 'pervasive' transcripts, such as antisense RNA species and promoter-upstream transcripts (PROMPTs), and of mRNAs with processing defects, thereby limiting or excluding their export to the cytoplasm. The RNA exosome may be involved in Ig class switch recombination (CSR) and/or Ig variable region somatic hypermutation (SHM) by targeting AICDA deamination activity to transcribed dsDNA substrates. In the cytoplasm, the RNA exosome complex is involved in general mRNA turnover and specifically degrades inherently unstable mRNAs containing AU-rich elements (AREs) within their 3' untranslated regions, and in RNA surveillance pathways, preventing translation of aberrant mRNAs. It seems to be involved in degradation of histone mRNA. The catalytic inactive RNA exosome core complex of 9 subunits (Exo-9) is proposed to play a pivotal role in the binding and presentation of RNA for ribonucleolysis, and to serve as a scaffold for the association with catalytic subunits and accessory proteins or complexes.[UniProtKB/Swiss-Prot Function]</p>


Product images:




Circular map for RC201419



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY EXOSC7 (Cat# RC201419, Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-EXOSC7 (Cat# [TA504239]). Positive lysates [LY402399] (100ug) and [LC402399] (20ug) can be purchased separately from OriGene.



Western blot validation of overexpression lysate (Cat# [LY402399]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC201419 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).



Coomassie blue staining of purified EXOSC7 protein (Cat# [TP301419]). The protein was produced from HEK293T cells transfected with EXOSC7 cDNA clone (Cat# RC201419) using MegaTran 2.0 (Cat# [TT210002]).