

Product datasheet for RC201454L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

RASD2 (NM_014310) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: RASD2 (NM 014310) Human Tagged ORF Clone Lentiviral Particle

Symbol: RASD2

Synonyms: Rhes; TEM2

Mammalian Cell Pur

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_014310

ORF Size: 798 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC201454).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 014310.3

 RefSeq Size:
 3047 bp

 RefSeq ORF:
 801 bp

 Locus ID:
 23551

 UniProt ID:
 Q96D21

 Cytogenetics:
 22q12.3

Domains: ras, RAN, RAS, RHO, RAB

Protein Families: Druggable Genome





ORIGENE

MW: 30.4 kDa

Gene Summary:

This gene belongs to the Ras superfamily of small GTPases and is enriched in the striatum. The encoded protein functions as an E3 ligase for attachment of small ubiquitin-like modifier (SUMO). This protein also binds to mutant huntingtin (mHtt), the protein mutated in Huntington disease (HD). Sumoylation of mHTT by this protein may cause degeneration of the striatum. The protein functions as an activator of mechanistic target of rapamycin 1 (mTOR1), which in turn plays a role in myelination, axon growth and regeneration. Reduced levels of mRNA expressed by this gene were found in HD patients. [provided by RefSeq, Jan 2016]