

Product datasheet for RC201462

PTRH2 (NM_016077) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: PTRH2 (NM_016077) Human Tagged ORF Clone
Tag: Myc-DDK
Symbol: PTRH2
Synonyms: BIT1; CFAP37; CGI-147; IMNEPD; PTH; PTH 2; PTH2
Mammalian Cell Selection: Neomycin
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
ORF Nucleotide Sequence: >RC201462 ORF sequence
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGCCCTCCAAATCCTTGGTTATGGAATATTTGGCTCATCCCAGTACACTCGGCTTGGCTGTTGGAGTTG
CTTGTGGCATGTGCCTGGGCTGGAGCCTTCGAGTATGCTTTGGGATGCTCCCAAAGCAAGACGAGCAA
GACACACACAGATACTGAAAGTGAAGCAAGCATCTTGGGAGACAGCGGGAGTACAAGATGATTCTTGTG
GTTCGAAATGACTTAAAGATGGGAAAAGGAAAGTGGTGCAGTCTCATGCTGCTGTTTCAGCCT
ACAAGCAGATTCAAAGAAGAAATCCTGAAATGCTCAAACAATGGGAATACTGTGGCCAGCCCAAGGTGGT
GGTCAAAGCTCCTGATGAAGAAACCTGATTGCATTATTGGCCCATGCAAAAATGCTGGGACTGACTGTA
AGTTTAATTCAAGATGCTGGACGTAAGTCAAGATTGCACCCAGGCTCTCAAATGTCCTAGGGATTGGCCAG
GACCAGCAGACCTAATTGACAAAGTCACTGGTCACCTAAAACCTTAC

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC201462 protein sequence
Red=Cloning site Green=Tags(s)

MPSKSLVMEYL AHPSTLGLAVGVACGMCLGWSLRVCFGMLPKSKTSKTHTDTESEASILGDSGEYKMILV
VRNDLKMGGKVAQAQC SHAAVSAYKQIQRRNPEMLKQWEYCGQPKVVVKAPDEETLIALLAHAKMLGLTV
SLIQDAGRTQIAPGSQTVLGIGP PADLIDKVTGHLKLY

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Chromatograms: https://cdn.origene.com/chromatograms/mk6300_f02.zip



[View online »](#)

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_016077

ORF Size: 537 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_016077.5](#)

RefSeq Size: 804 bp

RefSeq ORF: 540 bp

Locus ID: 51651

UniProt ID: [Q9Y3E5](#)


Cytogenetics: 17q23.1

Protein Families: Transmembrane


MW: 19.2 kDa

Gene Summary: The protein encoded by this gene is a mitochondrial protein with two putative domains, an N-terminal mitochondrial localization sequence, and a UPF0099 domain. In vitro assays suggest that this protein possesses peptidyl-tRNA hydrolase activity, to release the peptidyl moiety from tRNA, thereby preventing the accumulation of dissociated peptidyl-tRNA that could reduce the efficiency of translation. This protein also plays a role regulating cell survival and death. It promotes survival as part of an integrin-signaling pathway for cells attached to the extracellular matrix (ECM), but also promotes apoptosis in cells that have lost their attachment to the ECM, a process called anoikosis. After loss of cell attachment to the ECM, this protein is phosphorylated, is released from the mitochondria into the cytosol, and promotes caspase-independent apoptosis through interactions with transcriptional regulators. This gene has been implicated in the development and progression of tumors, and mutations in this gene have been associated with an infantile multisystem neurologic, endocrine, and pancreatic disease (INMEPD) characterized by intellectual disability, postnatal microcephaly, progressive cerebellar atrophy, hearing impairment, polyneuropathy, failure to thrive, and organ fibrosis with exocrine pancreas insufficiency (PMID: 25574476). Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Mar 2015]

Product images:



Circular map for RC201462



Western blot validation of overexpression lysate (Cat# [LY414202]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC201462 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).